

Part B. Section 1: Introduction

Since first published in 1980, the Dietary Guidelines for Americans have provided science-based advice to promote health and reduce risk of major chronic diseases through optimal diet and regular physical activity. The Dietary Guidelines have traditionally targeted the healthy general public older than age 2 years, but as data continue to accumulate regarding the importance of dietary intake during gestation and from birth on, it also will become important to consider those younger than age 2 years in future Guidelines. Because of their focus on health promotion and risk reduction, the Dietary Guidelines form the basis of Federal food, nutrition education, and information programs.

By law (Public Law 101-445, Title III, 7 U.S.C. 5301 et seq.) the most recent edition of the Dietary Guidelines is reviewed by a committee of experts, updated if necessary, and published every 5 years. The legislation also requires that the Secretaries of the US Department of Agriculture (USDA) and US Department of Health and Human Services (HHS) review all Federal publications for the general public containing dietary guidance information for consistency with the Dietary Guidelines for Americans. This Report presents the recommendations of the 2010 Dietary Guidelines Advisory Committee (DGAC) to the Secretaries of Agriculture and of Health and Human Services for use in updating the Guidelines.

The 2010 DGAC Report is unprecedented in addressing an American public, two-thirds of whom are overweight or obese. Americans are making dietary choices in a highly obesogenic environment and at a time of burgeoning diet-related chronic diseases affecting people of all ages, ethnic backgrounds, and socioeconomic levels. The DGAC considers the obesity epidemic to be the single greatest threat to public health in this century. This Report is therefore focused on evidence-based guidelines and recommendations that are considered effective and useful in halting and reversing the obesity problem through primary prevention and changes in behavior, the environment, and the food supply.

The Role of Diet and Physical Activity in Health Promotion: Attenuating Chronic Disease Risks

A large proportion of deaths each year in the US result from a limited number of preventable and modifiable factors. The leading causes of death for the past two decades have been tobacco use and poor diet and physical inactivity (McGinnis, 1993; Mokdad, 2004). The number of deaths related to poor diet and physical inactivity is increasing and may soon overtake tobacco as the leading cause of death. As discussed in this Report, poor dietary intake has been linked to excess body weight and numerous diseases and conditions, such as cardiovascular disease (CVD) and type 2 diabetes (T2D)

and their related risk factors. Even if the overweight/obesity epidemic resolves, the problems of chronic disease would continue to be a major health problems because poor-quality diets, even in the absence of overweight/obesity, increase the risk some of our most common chronic diseases.

The reduction of chronic disease risk merits strong emphasis in our Nation for many reasons, especially because some groups in the population bear a disproportionate burden of chronic disease and attendant risk factors. The present report highlights the evidence that links diet and the different chronic diseases. It also summarizes and synthesizes knowledge regarding many individual nutrients and food components into recommendations for an overall total pattern of eating that can be adopted by the public. Although adherence to the Dietary Guidelines is low among the US population, evidence is accumulating that selecting diets that comply with the Guidelines reduces the risk of chronic disease and promotes health. Ultimately, individuals choose the types and amount of food they eat and the amount of physical activity they perform, but the current environment significantly enhances the over-consumption of calories and discourages the expenditure of energy. Both sides of this equation are discussed in greater detail throughout the Report.

Population Groups of Particular Concern

The Dietary Guidelines for Americans has traditionally provided guidance to healthy Americans. However, the 2010 DGAC recognizes that a large percentage of the American population now has diet-related chronic diseases or risk factors for them, and has accommodated this reality in its review of the evidence. Much of the evidence the Committee reviewed pertains to adults. However, given the importance of nutrition across the lifespan and the rapidly growing scientific literature on diet and children's health, several sections of the Report focus particular attention on this important population group. In addition, the Committee presents reviews of evidence on several questions pertaining to pregnant and lactating women and to older adults.

Children

Increasingly, studies are addressing the role of nutrition and physical activity in promoting health in children. A nutrient-dense, high-quality diet, sufficient but not excessive in calories, and regular daily physical activity are integral to promoting the optimal health, growth, and development of children. For example, the rapid rates of growth occurring during adolescence increase the need for dietary sources of iron and calcium during that period to higher amounts per 1,000 calories than required at any other stage of life.

Evidence documents the importance of optimal nutrition starting during the fetal period through childhood and adolescence because this has a substantial influence on the risk of chronic

disease with age (Warner, 2010). Eating patterns established during childhood often are carried into adulthood (Aggett, 1994). For example, those who consume fruits and vegetables or milk regularly as children are more likely to do so as adults (Aggett, 1994).

Today, too many children are consuming diets with too many calories and not enough nutrients, and they are not getting enough physical activity (less than half of children age 12 to 21 years exercise on a daily basis [HHS, 1996]). As a result, chronic disease risk factors, such as glucose intolerance and hypertension, which were once unheard of in childhood, are now increasingly common. T2D now accounts for up to 50 percent of new cases of diabetes among youths. One in 400 youths will have T2D by age 20 years. Excess weight, particularly around the abdomen, as well as too little physical activity, appears to be the basis for developing this disease early in life.

Pregnant and Lactating Women

Both pregnancy and lactation are critical periods during which maternal nutrition is a key factor influencing the health of both child and mother. Energy as well as protein and several mineral and vitamin requirements increase substantially during pregnancy, making the pregnant woman's dietary choices critically important (Christian, 2010; IOM, 1991; IOM, 2002; Picciano, 2003).

However, excess energy intake during pregnancy has become a major concern. Growing evidence indicates that overnutrition leading to unhealthy weight gain during pregnancy may greatly predispose the child to obesity. Insufficient micronutrient intake also continues to be a concern. For example, sufficient intake of folic acid, which is especially important for normal development of the embryo and fetus, is critical during the entire periconceptional period. Dietary factors also may contribute to impaired glucose tolerance, a common disorder of pregnancy that influences fetal growth and outcomes (Clapp, 1998; Saldana, 2004). Dietary contaminants, such as methyl mercury, may adversely affect fetal growth. Maternal diet, especially the intake of certain vitamins and alcoholic beverages, also may influence breast milk composition (Dewey, 1999; IOM, 1991).

Older Adults

The *65+ in the United States: 2005 Report* noted that the US population aged 65 years and older is expected to double in size within the next 25 years (He, 2005). By 2030, it is projected that one in five people will be older than age 65 years. Individuals age 85 years and older are the fastest growing segment of the older population. In 2011, the "baby boom" generation will begin to turn 65. As the number of older Americans increases, the role of diet quality and physical activity in reducing the progression of chronic disease will become increasingly important. The health of older Americans is improving, but many are disabled and suffer from chronic conditions. The proportion with a

disability fell significantly from 26.2 percent in 1982 to 19.7 percent in 1999 (Manton, 2001), yet 14 million people age 65 years and older reported some level of disability in Census 2000, mostly linked to a high prevalence of chronic conditions, such as CVD, T2D, hypertension, or arthritis.

The process of aging can influence how nutrients are used and can exacerbate the effect of poor diet quality on health. For example, aging may reduce nutrient absorption, increase urinary nutrient loss, and alter normal pathways of nutrient metabolism. These changes associated with aging can be compensated to some extent by a nutrient-dense diet that remains within calorie needs. Most important, modifications of diet and increases in physical activity have tremendous potential as a means to prevent or delay chronic disease in older persons. Older individuals achieve, in many instances, greater benefit from a given improvement in diet than do younger individuals (e.g., older individuals tend to be more responsive to the blood pressure-lowering effects of reducing salt intake) or from an increase in physical activity. As with children, adolescents and younger adults, data comparing people aged 65 to 74 years in 1988-1994 and 1999-2000 show a startling rise in the percentage of obese older adults. In men, the proportion grew from about 24 to 33 percent and in women from about 27 percent to 39 percent (He, 2005).

Furthermore, available data have repeatedly documented that older-aged persons can make and sustain behavior change, more so than their younger counterparts (DPP, 2002; DPP, 2009; Whelton, 1997). Such results highlight the importance of encouraging dietary changes throughout the lifespan, including older-aged persons.

Changes in Diet and Physical Activity as a Means to Reduce Health Disparities

Of substantial concern are disparities in health among racial and ethnic minorities and among different socioeconomic groups. For example, Blacks have a higher prevalence of elevated blood pressure and a greater incidence of blood pressure-related diseases, such as stroke and kidney failure, than do non-Blacks (DGAC, 2004). Also, several subgroups of the population (e.g., Mexican-Americans, American Indians, and Blacks) have a strikingly high prevalence of overweight and obesity, even beyond that of the already high prevalence rates observed in the general population. Furthermore, it is well-recognized that individuals of lower socioeconomic status have a higher incidence of adverse health outcomes than do individuals of higher socioeconomic status. Dietary patterns differ among different groups, with individuals of lower education and income consuming fewer servings of vegetables and fruit than those with more education and higher income (USDA, 2004). While the reasons for such disparities are complex and multi-factorial, available research is sufficient to advocate certain dietary changes and increased physical activity as a means to reduce disparities.

The effects on blood pressure of a reduced sodium intake, increased potassium intake, and an overall healthy dietary pattern provide an example of how dietary changes could reduce health disparities. Although both Blacks and non-Blacks consume excess sodium, Blacks tend to be more sensitive to the effects of sodium than are non-Blacks. Likewise, Blacks tend to be more sensitive to the blood pressure-lowering effects of increased potassium intake. Ironically, the average potassium intake of Blacks is less than that of non-Blacks. The Dietary Approaches to Stop Hypertension (DASH) diet, an example of a healthy dietary pattern that emphasizes vegetables and fruits, has been shown in clinical trials to lower blood pressure to a greater extent in Blacks than in non-Blacks. Yet, Blacks tend to consume fewer fruits and vegetables than do non-Blacks.

Such evidence exemplifies important, yet underappreciated, opportunities to reduce health disparities through dietary changes.

From the 2010 DGAC Report to the Dietary Guidelines for Americans

A major goal of the 2010 DGAC is to summarize and synthesize the evidence to support USDA and HHS in developing nutrition recommendations that reduce the risk of chronic disease while meeting nutrient requirements and promoting health for all Americans.

The US Government uses the Dietary Guidelines as the basis of its food assistance programs, nutrition education efforts, and decisions about national health objectives. For example, the National School Lunch Program and the Elderly Nutrition Program incorporate the Dietary Guidelines in menu planning, the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) applies the Dietary Guidelines in its educational materials, and the Healthy People 2010 objectives for the Nation include objectives based on the Dietary Guidelines. The evidence described here in the 2010 DGAC Report, which will be used to develop the 2010 Dietary Guidelines for Americans, will help policymakers, educators, clinicians, and others speak with one voice on nutrition and health and to reduce the confusion caused by mixed messages in the media. The DGAC also hopes that the 2010 Dietary Guidelines for Americans will encourage the food industry to grow, manufacture, and sell foods that promote health and contribute to appropriate energy balance.

A Guide to the 2010 DGAC Report

This report contains several major components. Part A provides an Executive Summary to the Report. Part B sets the stage for the Report through this Introduction. It also provides a synthesis of major findings in two complementary chapters. The first chapter describes a health-promoting total

diet approach that combines the intake of foods, calories, and nutrients. The second chapter integrates the Report's major cross-cutting findings and provides specific recommendations for how Americans and different sectors throughout the Nation can put the Report's evidence-based dietary recommendations into action.

Part C describes the methodology the DGAC used to conduct its work and review the evidence on diet and health. Part D is the Science Base. In this Part, the DGAC's subcommittees present their specific findings in chapters focused on energy balance and weight management; nutrient adequacy; fatty acids and cholesterol; protein; carbohydrates; sodium, potassium, and water; alcohol, and food safety and technology.

The Report concludes with several Appendices, including a compilation of the Committee's scientific conclusions, a glossary, a brief history of the Dietary Guidelines for Americans, a listing of the food pattern analyses conducted for the 2010 DGAC, a summary of the process used to collect public comments, biographical sketches of DGAC members, and Acknowledgments.

References

- Aggett PJ, Haschke F, Heine W, Hernell O, Koletzko B, Lafeber H, Ormission A, Rey J, Tormo R. ESPGAN Committee on Nutrition Report: Childhood diet and prevention of coronary heart disease. *J Pediatr Gastr and Nutr* 1994;19(3):261-9.
- Clapp JF III. Effect of dietary carbohydrate on the glucose and insulin response to mixed caloric intake and exercise in both nonpregnant and pregnant women. *Diabetes Care* 1998;21(Suppl 2): B107-B112.
- Christian P. Micronutrients, birth weight, and survival. *Annu Rev Nutr* 2010 Apr 23; Epub ahead of print.
- Dewey KG, Schanler J, Koletzko B, eds. Nutrition and human lactation. *J Mammary Gland Biology & Neoplasia* 1999;4:241-95.
- Diabetes Prevention Program Research Group (DPP). The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 2002;25(12):2165-71.
- Diabetes Prevention Program Research Group, Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, Brenneman AT, Brown-Friday JO, Goldberg R, Venditti E, Nathan DM (DPP). 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;14;374(9702):1677-86.
- Dietary Guidelines Advisory Committee (DGAC). *Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2005*. Washington DC: US Department of Agriculture, Agricultural Research Service, August 2004.
- He W, Sengupta M, Velkoff V, DeBarros K. US Census Bureau. Current Population Reports. P23-209. *65+ in the United States: 2005*. Washington, DC: US Government Printing Office, 2005.
- Institute of Medicine. Subcommittee on Nutrition During Lactation. Committee on Nutritional Status During Pregnancy and Lactation. Food and Nutrition Board. *Nutrition During Lactation*. Washington, DC: National Academies Press, 1991.
- Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. Washington, DC: National Academies Press, 2002.
- Manton KG, Gu X. Changes in the prevalence of chronic disability in the United States black and nonblack population above age 65 from 1982 to 1999. *Proc Natl Acad Sci USA* 2001;98(11):6354-9.
- McCullough ML, Feskanich D, Stampher MJ, Giovannucci EL, Rimm EB, Hu FB, Spiegelman D, Hunter DJ, Colditz GA, Willett WC. Diet quality and major chronic disease risk in men and women: moving toward improved dietary guidance. *Am J Clin Nutr* 2002;76(6):1261-71.
- McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA* 1993;270(18):2207-12.

Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual Causes of Death in the United States, 2000. *JAMA* 2004; 291(10):1238-45. Correction: *JAMA* 2005;293(3):293-4.

Picciano MF. Pregnancy and lactation: physiological adjustments, nutritional requirements and the role of dietary supplements. *J Nutr* 2003 Jun;133(6):1997S-2002S.

Saldana TM, Siega-Riz AM, Adair LS. Effect of macronutrient intake on the development of glucose intolerance during pregnancy. *Am J Clin Nutr* 2004;79(3):479-86.

US Department of Agriculture (USDA). Continuing Survey of Food Intakes by Individuals 1994-1996, 1998. PB2000-500027. CD-ROM, 2004.

US Department of Health and Human Services (HHS). *Physical Activity and Health: A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention. National Center for Disease Prevention and Health Promotion, 1996.

Warner MJ, Ozanne SE. Mechanisms involved in the developmental programming of adulthood disease. *Biochem J* 2010 Apr 14;427(3):333-47.

Part B. Section 2: The Total Diet: Combining Nutrients, Consuming Food

Introduction

The 2010 Dietary Guidelines Advisory Committee (DGAC) supports a total diet approach to achieving dietary goals. The purpose of this chapter is to demonstrate how the scientific evidence presented in each of the topic-specific chapters in *Part D: The Science Base*—Energy Balance and Weight Management; Nutrient Adequacy; Fatty Acids and Cholesterol; Protein; Carbohydrates; Sodium, Potassium, and Water; Alcohol; and Food Safety and Technology—can be incorporated into an overall eating pattern that optimizes health outcomes.

Until recently, data were insufficient to document the impact of whole diets and eating patterns on health outcomes. The state of the evidence and the methodologic rigor regarding such questions have improved tremendously and the data can now be incorporated into this Report.

This chapter synthesizes the evidence on dietary components that contribute to excess energy and inadequate nutrient intakes in the United States (US), and the foods that can provide these missing essential nutrients and other health benefits. It presents a brief, evidence-based comparison of worldwide eating patterns, including the Dietary Approaches to Stop Hypertension (DASH), Mediterranean, and other patterns, along with a description of the US Department of Agriculture (USDA) Food Patterns with vegetarian variations.

A nutrient-dense total diet has multiple health benefits and can be implemented in various ways. The US is comprised of individuals of all ages who come from many cultures and have a variety of food and taste preferences. All of these factors were considered in developing a recommended total diet that is flexible while meeting nutrient needs without exceeding energy requirements.

The Catalyst for the Total Diet Approach

Although there is no single “American” or “Western” diet, average American food patterns currently bear little resemblance to the diet recommended in the 2005 Dietary Guidelines for Americans. As documented by the latest data from the National Health and Nutrition Examination Survey (NHANES), Americans eat too many calories and too much solid fats, added sugars, refined grains, and sodium. Americans also eat too little dietary fiber, vitamin D, calcium, potassium, and unsaturated fatty acids (specifically omega-3s), and other important nutrients that are mostly found in vegetables, fruits, whole grains, low-fat milk and milk products, and seafood (see *Part D. Section 2: Nutrient Adequacy*).

Overweight and obesity are highly prevalent in the US in both adults and children. This is of great public health concern because excess body fat is associated with a much higher risk of premature death and many serious disorders, as identified in *Part D. Section 1: Energy Balance and Weight Management*. Preventing overweight is highly preferable to initiating weight loss treatment after weight gain occurs, because the failure rate in achieving and maintaining weight loss is very high. Furthermore, the behaviors required to prevent overweight are less daunting than the behaviors necessary to lose and sustain weight loss. Currently, the average American gains about a pound a year between the ages of 20 to 60 years. Some persons gain much more. Remaining conscious of one's body weight throughout life and adopting a lifestyle early on that will achieve and sustain weight control across the lifespan are paramount to maintaining good health and quality of life.

A Special Focus on Children and Adolescents

The single most significant adverse health trend among US children in the past 40 years has been the dramatic increase in overweight and obesity (see *Part D. Section 1: Energy Balance and Weight Management*). Since the early 1970s, the prevalence of overweight and obesity has approximately doubled among children ages 2 to 11 years, and tripled among adolescents ages 12 to 19 years. Not only is obesity associated with adverse health effects during childhood, but evidence documents increased risk of future chronic disease in adult life.

Childhood obesity results from poorly regulated energy balance. Ideally, children and adolescents should consume foods that provide an adequate intake of all essential nutrients needed for normal growth and development, metabolism, immunity and cognitive function, without exceeding caloric requirements. Factors associated with preventing excess adiposity in children are incorporated into the total diet described here, and include:

- Energy intake balanced with expenditure
- Greatly reduced intake of sugar-sweetened beverages
- Increased intake of vegetables and fruits
- Smaller amounts of fruit juice, especially for overweight children
- Smaller portions of foods and beverages
- Infrequent consumption of meals from quick service (i.e., fast food) restaurants
- Habitual consumption of breakfast
- Fewer hours of screen time (e.g., television, computer)
- More hours of active play

Blending Science-Based Recommendations Into a Healthful Total Diet

The DGAC defines “total diet” as the combination of foods and beverages that provide energy and nutrients and constitute an individual’s complete dietary intake, on average, over time. This encompasses various foods and food groups, their recommended amounts and frequency, and the resulting eating pattern. To achieve dietary goals and energy balance, Americans must become mindful, or “conscious,” eaters, that is, attentively choosing what and how much they eat. Since the mid-1980s, the USDA has provided recommended food patterns that represent a total diet approach to dietary guidance (Britten, 2006). The most recent USDA Food Patterns have been visually conveyed as the MyPyramid Food Guidance System (Haven, 2006). This approach was intended to help people personalize dietary recommendations and offer flexibility based on individual preferences. The key core components of a nutrient-dense total diet for all Americans are presented below.

Moderate Energy Intake

The DGAC encourages Americans to achieve their recommended nutrient intakes by consuming foods within a total diet that meets but does not exceed energy needs. Overweight and obesity result from energy imbalance (intake exceeding expenditure) (see *Part D. Section 1: Energy Balance and Weight Management*). The increased incidence and current high proportion of overweight and obesity in the US illustrates an energy imbalance across virtually all subgroups of the population. People consume too many calories (i.e., energy) relative to the calories they expend. As a start, all Americans are encouraged to know their energy needs in order to avoid inappropriate weight gain. Table B2.1 (see the end of this chapter) can help individuals identify their energy needs based on their age, sex, and level of activity. Self-monitoring of both calorie intake and time spent in physical activity is one of the most useful tools a person can use to engage in and maintain behaviors that sustain a healthy weight.

Because levels of leisure time physical activity in US adults have remained stable or increased only slightly between 1990 and 2004, it is clear that an increased calorie intake has been the primary cause of the obesity problem. Hence, even though one can achieve a calorie deficit by increasing physical activity, the primary focus should be on reducing excessive calorie intake.

Overall, the top food sources of energy, and mean energy intake from each, for the US population, as reported in the National Health and Nutrition Examination Survey (NHANES) 2005-2006, are (NCI, 2010a):

- Grain-based desserts (cakes, cookies, doughnuts, pies, crisps, cobblers, and granola bars; 139 calories per day)

- Yeast breads (129 calories per day)
- Chicken and chicken mixed dishes (121 calories per day)
- Soda/energy/sports drinks (114 calories per day)
- Pizza (98 calories per day)

While the top sources of energy intake vary by age group, many of these sources are foods and beverages that are not in nutrient-dense forms. For example, the top energy source for adults ages 19 years and older and for children ages 4 to 13 years is grain-based desserts. These desserts are also among the top five sources of energy for teens and younger children. For teens ages 14 to 18 years, the top energy source is soda/energy/sports drinks, and these beverages are also among the top five energy sources for adults ages 19 years and older and for children ages 9 to 13 years. For children ages 2 to 3 years only, the top energy source is whole milk (rather than low-fat milk). Other foods that are among the top five sources of energy for various age groups are yeast breads, chicken and chicken mixed dishes, pizza, and, for adults only, alcoholic beverages (NCI, 2010a; see Table B2.2 at the end of this chapter for the top five sources of energy for each age group, and Tables D1.1, D1.6, and D1.7 in *Part D. Section 1: Energy Balance and Weight Management* for more detailed lists of food sources of energy).

Total diets that are high in energy, but low in nutrients, can paradoxically leave a person overweight but undernourished and thus, at higher risk of cardiovascular disease (CVD), type 2 diabetes (T2D), and certain types of cancers. Of urgent concern is America's youth, most of whom currently fit this pattern. Many children consume nutrient-poor sources of energy at the highest end of their respective energy ranges (see Figure D1.1 in *Part D. Section 1: Energy Balance and Weight Management*) and they are increasingly sedentary.

Beverages also contribute substantially to overall dietary and energy intake. Although they provide needed fluid, beverages often add calories to the diet without providing nutrients. Their consumption should be planned in the context of total calorie intake and how they can fit into the total diet of each individual. Currently, US adults ages 19 years and older consume an average of 394 calories per day as beverages. The major types of beverages consumed, and the mean caloric intake from each, are (NCI, 2010b):

- Soda (114 calories per day)
- Coffee and tea (26 calories per day)
- Fluid milk (80 calories per day)
- 100 percent fruit juice and fruit drinks (67 calories per day)

- Alcoholic beverages (108 calories per day)

Children, ages 2 to 18 years, consume an average of 400 calories per day as beverages. The major beverages for children and calories from each are somewhat different:

- Fluid milk (162 calories per day)
- Soda (121 calories per day)
- 100 percent fruit juices and fruit drinks (112 calories per day)

In children, the amount and source of calories from beverages differs by age. For example, 100 percent fruit juice is a prominent source of energy in children ages 2 to 3 years, while soda/sports/energy drinks are the most common source of energy among beverages (and energy overall) in children ages 14 to 18 years.

Portion control and the quantity of foods and beverages consumed within the total diet also are important considerations in moderating energy intake (see *Part D. Section 1: Energy Balance and Weight Management*). Excessive portion sizes are very common in the US and are linked to higher energy intakes and weight gain over time. This is particularly true when large portions of foods high in solid fats and added sugars (SoFAS) and refined grains are consumed.

Reduce Solid Fats and Added Sugars (SoFAS)

Solid fats and added sugars contribute substantially (approximately 35% of calories) to total energy intakes of Americans, thereby leading to excessive saturated fat and cholesterol intakes and insufficient intake of dietary fiber and other nutrients (see *Part D. Section 2: Nutrient Adequacy*; *Part D. Section 3: Fatty Acids and Cholesterol*; and *Part D. Section 5: Carbohydrates*).


The 2005 DGAC defined the term “discretionary calorie allowance” as “the difference between total energy requirements and the energy consumed to meet recommended nutrient intakes” (DGAC, 2004). Discretionary calories were intended to represent the calories available for consumption *only* after meeting nutrient recommendations and without exceeding total energy needs. Unfortunately, this concept has been difficult to translate into meaningful consumer education. To clarify translation, the 2010 DGAC focused specifically on reducing the intake of solid fats and added sugars (SoFAS), which provide most of the non-essential or extra calories that Americans consume. Major food sources of the two components of SoFAS are (Bachman, 2008):

- Solid fats (percent of solid fat intake)
 - Grain-based desserts, including cakes, cookies, pies, doughnuts, and granola bars (10.9%)
 - Regular cheese (7.7%)
 - Sausage, franks, bacon, and ribs (7.1%)

- Pizza (5.9%)
- Fried white potatoes, including French fries and hash browns (5.5%)
- Dairy-based desserts, such as ice cream (5.1%)
- Added sugars (percent of added sugars intake)
 - Soda (36.6%)
 - Grain-based desserts (11.7%)
 - Fruit drinks (11.5%)
 - Dairy-based desserts (6.4%)
 - Candy (6.2%)

Maximum limits on SoFAS are meant to be estimates and not necessarily daily targets (see limits from USDA Food Patterns, Table B2.3, end of this chapter). These foods should constitute a very small proportion of total energy intake in the total diet. Figure B2.1 contrasts the current disproportionately high intake of SoFAS with what is more appropriate from a healthy eating pattern.

Figure B2.1. What we eat versus recommended limits: Calories from Solid Fats and Added Sugars (SoFAS)



Note: The depiction of the proportionate amounts of total calories consumed and the recommended limits are illustrative only. The figure illustrates about 35 percent of total calories consumed as SoFAS, on average, in contrast to a recommended limit of no more than about 5 to 15 percent of total calories for most individuals.


Americans currently consume 35 percent of their total calories from SoFAS. This is too high. They should reduce intake of calories from SoFAS by 20 to 30 percent. This means that no more than 5 to 15 percent of total calories should be derived from SoFAS. For example, the USDA Food Patterns limit SoFAS to about 120 calories in the 1600-calorie pattern, 160 calories in the 1800-calorie pattern, and 260 calories in the 2000-calorie pattern (Table B2.3, at the end of the chapter, lists SoFAS limits for all calorie levels). Reduction of calories from SoFAS to these amounts allows

for increased intakes of nutrient-dense foods such as vegetables (including cooked dry beans and peas), fruits, whole grains, and fat-free and low-fat fluid milk and milk products, without exceeding overall calorie needs.

Consume Nutrient-Dense Foods (But Not too Much of Them)

Currently, Americans consume less than 20 percent of the recommended intakes for whole grains, less than 60 percent for vegetables, less than 50 percent for fruits, and less than 60 percent for milk and milk products (Figure B2.2). Inadequate intakes of nutrient-dense foods from these basic food groups place individuals at risk for lower than recommended levels of specific nutrients, namely vitamin D, calcium, potassium, and dietary fiber.

Figure B2.2. Dietary intakes in comparison to recommended intake levels or limits



Note: Bars show average intakes for all individuals (ages 1 or 2 years or older) as a percent of the recommended intake level or limit. Recommended intakes for food groups and limits for refined grains, SoFAS, solid fats, and added sugars are based on the USDA 2000-calorie food patterns. Recommended intakes for fiber, potassium, vitamin D, and calcium are based on the highest AI for ages 14 to 70 years. Limits for sodium are based on the AI and for saturated fat on 7 percent of calories.

Data source: What We Eat in America, National Health and Nutrition Examination Survey (WWEIA, NHANES) 2001-2004 or 2005-2006.

Figure B2.2. Data points. All values in percents.

	Percent of recommended intake
Whole grains	15
Vegetables	59
Fruits	42
Milk	52
Oils	61
Fiber	40
Potassium	56
Vitamin D	42
Calcium	75
	Percent of recommended limit
Calories from SoFAS	280
Added sugars	242
Solid fats	281
Refined grains	200
Sodium	229
Saturated fat	158

Food from all food groups are composed of a combination of the macronutrients carbohydrates, fats, and protein in varying proportions. These are the major sources of energy in any food or diet. Understanding their role in the diet will help Americans make appropriate food choices.

Carbohydrates (4 kcal/g) are the primary source of energy intake, and higher intakes of carbohydrates, especially complex sources, are recommended for active people. Sedentary individuals, and thus most Americans, should lower their intakes of refined carbohydrates, greatly reducing intakes of sugar and sugar-sweetened beverages and refined grains that are high in calories, but relatively low in certain nutrients. Whole-grain versions of many grain products (such as plain white bread, rolls, bagels, muffins, pasta, breakfast cereals) should be substituted to meet the recommendation that half of grains consumed be whole grains, also assisting in meeting dietary fiber recommendations (see *Part D. Section 5: Carbohydrates*).

Dietary fats (both solid fats and oils) are high in calories (9 kcal/g). Unsaturated fats, including omega-3 from seafood sources, should be increased and saturated fat and *trans* fatty acid intake should be minimized. Given typical patterns of consumption in the US, dietary saturated fat intake is highly correlated with total fat intake. Consuming the recommended intake of saturated fat (less than 10% of calories immediately as an interim step toward an eventual goal of less than 7% of calories)

is more likely achievable when total fat intake is less than 30 percent of total calories. It is recommended that total fat should be in the range of 20 to 35 percent of total calories but derived mostly from oils within a nutrient-rich, energy-balanced dietary pattern. These oils should replace solid fats and not add calories to the total diet (see *Part D. Section 3: Fatty Acids and Cholesterol*).

Dietary protein (4 kcal/g) provides essential amino acids and energy, and assists in building and preserving body proteins. Both plant-based sources of protein (i.e., cooked dry beans and peas, nuts, seeds, and soy products) and animal-based sources (i.e., meat, poultry, seafood, eggs, and low-fat and fat-free milk) can be incorporated into the total diet, with further emphasis on increasing seafood (rich in omega-3 fatty acids as well as protein) and cooked dry beans and peas (rich in dietary fiber as well as vegetable protein) (see *Part D. Section 4: Protein*).

Consumption of alcoholic beverages also contributes to calories (7 kcal/g), from the alcohol itself as well as accompanying mixers (e.g., soda, juice or sweetened mixer). In many cases, the accompanying mixer (See Table D1.9 in *Part D. Section 1: Energy Balance and Weight Management*) has more calories than the alcohol itself, so careful attention to portion size is important for alcoholic beverages. Based on individual preferences among adults, a moderate amount of alcohol may be included in the total diet if calorie allowances are not exceeded and essential nutrient needs are met. For adults who are attempting to reduce calorie intake, alcohol could be one of the energy sources that is reduced to lower total calorie intake. Pregnant women or individuals with certain medical conditions or on certain medications as well as individuals who will take part in activities that require attention or skill should not consume alcohol (see *Part D. Section 7: Alcohol*).

Vegetables, fruits, high-fiber whole grains, seafood, eggs, and nuts prepared without added SoFAS are considered “nutrient-dense foods,” as are low-fat forms of milk and lean meat and poultry prepared without added SoFAS. Nutrient-dense foods are found in a variety of forms but ideally are minimally processed and minimize or exclude added SoFAS, starches, and sodium. Combined into a total diet, these foods should provide a full range of essential nutrients, including those of special concern (e.g., vitamin D, calcium, potassium, and dietary fiber).

Finally, the nutrient-dense total diet should be prepared using best practices for food safety to ensure that foods consumed do not induce foodborne illnesses. (See *Part D. Section 8: Food Safety and Technology*.) A balanced grouping of a variety of foods among all the food groups, consumed in moderation, that are culturally appealing will offer pleasurable eating experiences and promote health among Americans.

Reduce Sodium Intake

Even a nutrient-dense total diet that remains excessive in sodium can lead to health consequences such as elevated blood pressure. Excessive sodium intake raises blood pressure, a well-documented and extraordinarily common risk factor for heart disease, stroke, and kidney disease. Although most research has been conducted in adults, the adverse effects of sodium on blood pressure begin early in life, and reducing sodium intake has substantial health benefits. Given the fact that a higher potassium intake attenuates the adverse effects of sodium on blood pressure, ensuring increased intakes of dietary potassium also would have health benefits.

The current food supply is replete with excess sodium. In this setting, virtually all Americans exceed the recommended upper limit of sodium intake. Because approximately 75 percent of dietary sodium is added during food processing, food manufacturers and restaurant industries have a critically important role in reducing the sodium intake. In addition, individuals should choose and prepare foods with little or no sodium (see *Part D. Section 6: Sodium, Potassium, and Water*).

A Flexible Approach to Applying Total Diet Recommendations

A healthful total diet is not a rigid prescription, but rather is a flexible approach that incorporates a wide range of individual tastes and preferences. Just as there is no one “American” or “Western” diet, there is no one recommendation for a healthful diet. As is evident in the following sections, data are accumulating that certain dietary patterns consumed around the world are associated with beneficial health outcomes. Likewise, the Food Patterns developed by the USDA illustrate that both nutrient and moderation goals can be met in a variety of ways.

Worldwide Dietary Patterns Provide Support for a Nutrient-dense Total Diet

Across the world and within the US, there are striking differences in diets and also in diet-related health outcomes. Although research on dietary patterns is complex, and many methodological issues remain in synthesizing data across studies, a consensus is emerging that consumption of certain dietary patterns is associated with a reduced risk of several major chronic diseases. The 2010 DGAC focused on the effects of dietary patterns on total mortality, CVD, and blood pressure (a major diet-related cardiovascular risk factor). The World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR, 2007) recently reviewed the available evidence of the relationship of cancer with specific dietary factors and overall dietary patterns. While several dietary factors were associated with specific types of cancer, it concluded that no firm judgment can be made on the relationship of dietary patterns with cancer.

The 2010 DGAC focused on the DASH-style dietary patterns and Mediterranean-style dietary patterns because considerable research exists on health outcomes as well as information on nutrient and food group composition. It also examined traditional Asian dietary patterns and vegetarian diets. Traditional Asian dietary patterns (e.g., Japanese and Okinawan dietary patterns) have been associated with a reduced risk of coronary heart disease, but documentation using contemporary research methods is scant. Most traditional dietary patterns provide for health at least moderately well, and their variety demonstrates that a person can eat healthfully in a number of ways. Vegetarian diets have been associated with a reduced risk of CVD, but information on nutrient content and food group composition is sparse.

Dietary patterns with health benefits are summarized below. An Appendix at the end of this chapter provides further detail on these dietary patterns as well as several summary tables.

DASH-style Dietary Patterns

DASH-style dietary patterns emphasize vegetables, fruits, and low-fat milk and milk products; include whole grains, poultry, seafood, and nuts; and are reduced in red meat, sweets, sodium, and sugar-containing beverages. As originally tested, the DASH diet is reduced in total fat (27% of kcal) with total protein intake of 18 percent of calories and carbohydrate intake of 55 percent of calories. However, other versions of the DASH diet are available, in which carbohydrate is partially replaced with protein (about half from plant sources) or unsaturated fat (predominantly monounsaturated fat). The latter version is noteworthy because nutrient adequacy and a reduced saturated fat intake (6% of kcal) were both achieved in the setting of high monounsaturated fat (21% of kcal) and total fat (37% of kcal) intake. In a free-living setting, care is needed to meet but not exceed energy needs in order to avoid weight gain.

Each of these DASH style diets lowers blood pressure, improves blood lipids, and reduces CVD risk. Blood pressure reduction is the greatest when the DASH diet is consumed with reduced sodium intake. At present, few adults, even those with hypertension, eat a diet that is consistent with the DASH dietary pattern.

Mediterranean-style Dietary Patterns

In view of the large number of cultures and agricultural patterns of countries that border the Mediterranean Sea, the “Mediterranean” diet is not a single dietary pattern. Although no well-accepted set of criteria exist, a traditional Mediterranean diet can be described as one that emphasizes breads and other cereal foods usually made from wheat, vegetables, fruits, nuts, unrefined cereals, and olive oil; includes fish and wine with meals (in non-Islamic countries); and is reduced in saturated fat, meat, and full-fat dairy products. Results from observational studies and

clinical trials suggest that consumption of a traditional Mediterranean diet, similar to that of Crete in the 1960s, is associated with one of the lowest risks of coronary heart disease in the world. Over time, the diet of Crete has changed remarkably and is now characterized by higher intake of saturated fat and cholesterol, and reduced intake of monounsaturated fats. At the same time, total fat consumption has fallen. These trends have been accompanied by a steady rise in heart disease risk.

Vegetarian Dietary Patterns

In some observational studies, vegetarian diets and lifestyle have been associated with improved health outcomes. The types of vegetarian diets consumed in the US vary widely. Vegans do not consume any animal products, while lacto-ovo vegetarians, consume milk and eggs. Although not strict vegetarians, many individuals consume small or minimal amounts of animal products. On average, vegetarians consume fewer calories from fat than non-vegetarians, particularly saturated fat, and have a higher consumption of carbohydrates than non-vegetarians. In addition, vegetarians tend to consume fewer overall calories and have a lower body mass index than non-vegetarians. These characteristics, in addition to the dietary pattern per se, may contribute to the improved health outcomes of vegetarians (see the Appendix at the end of this chapter and ***Part D. Section 4: Protein*** for additional information on vegetarian diets).

Other Dietary Patterns

In view of the increasing diversity of the US population, interest in the health effects of non-Western diets is substantial. One group of diets with potential health benefits are those traditionally consumed in Asia, which has experienced some of the lowest rates of coronary heart disease in the world. Both traditional Japanese and Okinawan dietary patterns have been associated with a low risk of coronary heart disease. Nonetheless, compared to the evidence supporting DASH and Mediterranean diets, detailed information on diet composition as well as epidemiologic and clinical trial evidence on health benefits, similar to that available for the other types of diets, is sparse. Also, over time, dietary intakes in these countries have changed and may no longer reflect the healthiest choices.

USDA Food Patterns Provide Guidance for Meeting Dietary Guideline Recommendations

Applying results from carefully conducted studies of nutrition and health, the USDA has developed a number of different food guides over the past century. These guides have identified eating patterns that meet known nutrient needs and balance intake from various food groups. Based

upon the Nation’s dietary intake at the time, early USDA food guides focused on nutrient adequacy only. Due to the health risks associated with overconsumption of specific dietary components, including the increasing obesity problem, recent guides have encompassed moderation goals while meeting nutrient adequacy goals. The current USDA Food Patterns also are aimed at primary disease prevention. For example, Table B2.4 (see end of chapter) compares the 2000-calorie USDA food pattern with the DASH diet and with current consumption patterns. The types and amounts of foods recommended in the USDA patterns are very similar to the DASH diet, and both are very different from current intakes.

The USDA Food Patterns recommend the amounts of foods to eat each day from the five major food groups and subgroups, specifically in nutrient-dense forms. The Patterns allow for oils and limit the maximum number of calories that should be consumed from SoFAS. Table B2.3 (see end of chapter) shows recommended amounts and limits in the USDA food patterns at all 12 energy levels (*Part D. Section 2: Nutrient Adequacy*, Table D2.1 provides the specific nutritional goals for each pattern).

The USDA Food Patterns incorporate several important assumptions:

- A variety of foods are used to meet recommended intakes from each food group or subgroup, in amounts proportionate to current consumption by the population.
- Food choices selected for use in the analysis are in nutrient-dense forms, that is, with little or no SoFAS, and in most cases without added salt.
- For each age-sex group, the pattern developed to meet nutrient needs is at a caloric level that meets but does not exceed energy needs for sedentary individuals.

The online *Appendix E3.1: Adequacy of the USDA Food Patterns*, available at www.dietaryguidelines.gov, provides details of the analysis conducted for the DGAC to determine whether the USDA Food Patterns meet nutritional goals for adequacy and moderation while staying within established calorie targets.

Recommended intake amounts in the USDA Food Patterns remain unchanged from 2005 with the exception of the vegetable subgroups. Several changes were made to decrease the wide discrepancy in number and amounts of vegetables consumed between the largest and the smallest subgroups. This resulted in moving tomatoes and red peppers from “other vegetables” to a new “red-orange vegetable” subgroup, which provided a greater focus on tomatoes without compromising the nutrient adequacy of the patterns (see the online *Appendix E3.2: Realigning Vegetable Subgroups* report at www.dietaryguidelines.gov, for details). The USDA Food Patterns meet almost all of their nutritional goals for adequacy and moderation, when evaluated using current food composition and consumption data.

USDA also developed and evaluated several variations on the base patterns, applying the same principles but modifying food choices to accommodate those wanting to eat a plant-based or vegetarian diet. An additional analysis investigated a possible modification of the patterns for those tracking carbohydrate intake, such as people with diabetes. The results of these analyses are presented below (see *Part C: Methodology* for a description of the methods used and a list of all food pattern modeling analyses).

Vegetarian Patterns Based on USDA Food Patterns

The USDA Food Patterns include two animal-based food groups: the “meat, poultry, seafood, eggs, soy products, nuts, and seeds” group and the “milk, yogurt, and cheese” group. Although the groups contain some plant foods, the majority of consumption from them is from animal products. As is true in American diets, these two food groups in the Food Patterns are the major sources of protein, calcium, vitamin D, vitamin B₁₂, riboflavin, choline, selenium, zinc, and the omega-3 fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA).

The USDA food patterns were modified to replace some or all animal products with plant products (see the online *Appendix E3.3: Vegetarian Food Patterns* report at www.dietaryguidelines.gov for details). The plant-based (at least 50% of all protein from plant sources), lacto-ovo vegetarian (no meat, poultry, or seafood), and vegan (no meat, poultry, seafood, eggs, fluid milk or milk products) food patterns, collectively referred to as the “vegetarian patterns,” meet almost all goals for nutrient adequacy. Amounts of protein, including all essential amino acids, were adequate in all vegetarian patterns. Amounts of calcium and vitamins D and B₁₂ were adequate because fortified sources of these nutrients were selected to replace milk and meat products. The estimated bioavailable iron in the vegan patterns was less than the RDA for some children and women. While no dietary standards exist for omega-3 fatty acids, levels of EPA and DHA are substantially lower than the base Food Patterns, especially in the vegan patterns. All moderation goals are met in the vegetarian patterns. If only plant foods are consumed, choices should include foods fortified with vitamin B₁₂, vitamin D, and calcium. Other nutrients of potential concern include iron, choline, EPA, and DHA.

Considering an Alternative Placement for Starchy Vegetables

To offer flexibility in selecting a food pattern that meets nutrient needs and accommodates food preferences, USDA evaluated a nutritionally adequate option that considers starchy vegetables as a grain alternative (See the online *Appendix E3.4: Starchy Vegetables* report at www.dietaryguidelines.gov for details). This pattern may be useful for individuals who wish to track the amount of carbohydrates they consume, who prefer a dietary pattern that groups all major

sources of starch together, or who wish to integrate the USDA recommendations with other diet plans. In this pattern, individuals can substitute starchy vegetables for a portion of the recommended grains, as long as they eat additional vegetables from other subgroups to replace the starchy vegetables. As with all of the modeling analyses, the vegetables and grains selected should be nutrient-dense forms, not forms with added fats, sugars, or salt. Although starchy vegetables remain part of the vegetable group in the USDA Food Patterns, this analysis identified an option for flexibility to help some individuals integrate the USDA recommendations with other dietary plans.

The Importance of Nutrient-dense Choices

The USDA Food Patterns assume that foods in each food group will be consumed in the same relative proportions as they appear in the average American diet, but that most will be in nutrient-dense forms. Nutrient-dense choices are available to consumers, but they are not the forms most typically consumed. Consuming recommended amounts of foods, but in forms that represent typical food choices rather than the “ideal” nutrient-dense choices, has a major impact on energy and nutrient intake. Excess intake of energy, sodium, saturated fat, and cholesterol result from using typical food choices in the recommended amounts for the patterns. For example, assuming typical food choices, the calorie intake in the 2000-calorie pattern is almost 400 calories more per day than the target (see the online *Appendix E3.5: “Typical Choices” Food Patterns* report at www.dietaryguidelines.gov for details of an analysis of the effect of typical versus ideal choices). If consumers act on the message about quantities to eat from each food group or subgroup, but fail to implement the moderation messages about choosing most foods in low-fat, no-added-sugars, and low-sodium forms, they will not meet the important moderation goals.

Chapter Summary

Good health and vitality across the life span are what Americans desire. The 2010 DGAC report concludes that this is achievable but requires a lifestyle approach that includes a total diet that is:

- Energy balanced, limited in total calories, and portion controlled
- Nutrient-dense and includes:
 - Vegetables, fruits, high-fiber whole grains
 - Fat-free or low-fat fluid milk and milk products
 - Seafood, lean meat and poultry, eggs, soy products, nuts, seeds, and oils
- Very low in solid fats and added sugars (SoFAS)

- Reduced in sodium

Physical activity will assist in the helping to achieve a balance between calorie intake and expenditure, leading to body weight maintenance. Children and adolescents are of particular concern because the dietary habits that they form during their youth will set the foundation for their choices and behaviors as adults.

Several distinct dietary patterns are associated with health benefits, including lower blood pressure and a reduced risk of CVD and total mortality. A common feature of these diets is an emphasis on plant foods. Accordingly, fiber intake is high and saturated fat is typically low. When total fat intake is high, that is, more than 30 percent of calories, the predominant fats are monounsaturated and polyunsaturated fats. Carbohydrate intake is typically in the range of 50 to 60 percent of calories, but these often include whole grain products with minimal processing, as well as cooked dry beans and peas. The totality of evidence documenting a beneficial impact of plant-based dietary patterns on CVD risk is remarkable and worthy of recommendation.

Americans have considerable flexibility in selecting a diet that includes foods they enjoy, meets nutrient requirements, reduces risk of preventable disease, and controls weight. No one specific dietary pattern provides the only way to incorporate the principles listed above into a total diet. The daunting public health challenge is to accomplish population-wide adoption of healthful dietary patterns within the setting of powerful influences that currently promote unhealthy lifestyles. The 2010 DGAC is united in advocating that policy makers, stakeholders and health-care providers embrace and support these important, evidence-based guidelines for the benefit of all Americans.

References

Bachman JL, Reedy J, Subar AF, Krebs-Smith SM. Sources of food group intakes among the US population, 2001-2002. *J Am Diet Assoc.* 2008;108(5):804-14.

Britten P, Marcoe K, Yamini S, Davis C. Development of food intake patterns for the MyPyramid Food Guidance System. *J Nutr Educ Behav.* 2006;38(6 Suppl):S78-S92.

Dietary Guidelines Advisory Committee (DGAC). *Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2005.* Washington DC: US Department of Agriculture, Agricultural Research Service, August 2004.

Haven J, Burns A, Britten P, Davis C. Developing the consumer interface for the MyPyramid Food Guidance System. *J Nutr Educ Behav.* 2006;38(6 Suppl):S124-S135.

Marcoe K, Juan W, Yamini S, Carlson A, Britten P. Development of food group composites and nutrient profiles for the MyPyramid Food Guidance System. *J Nutr Educ Behav.* 2006;38(6 Suppl):S93-S107.

National Cancer Institute (NCI). Food Sources of Energy Among US Children and Adolescents, 2005-2006. Risk Factor Monitoring and Methods Branch Website. Applied Research Program. National Cancer Institute, 2010a. <http://riskfactor.cancer.gov/diet/foodsources/>. Updated May 21, 2010. Accessed May 21, 2010.

National Cancer Institute (NCI). Distribution of Intake across Beverage Types, US Population, 2005-2006. Risk Factor Monitoring and Methods Branch Website. Applied Research Program. National Cancer Institute, 2010b. <http://riskfactor.cancer.gov/diet/foodsources/>. Updated May 21, 2010. Accessed May 21, 2010.

World Cancer Research Fund and American Institute for Cancer Research Report (WCRF/AICR). *Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective.* Washington, DC: AICR, 2007.

Table B2.1. Estimated energy needs¹ in calories per day, for reference-sized individuals by age, sex, and activity level

Sex/Activity level	Male/ Sedentary	Male/ Moderately Active	Male/ Active	Female/ Sedentary	Female/ Moderately Active	Female/ Active
Age						
2	1000	1000	1000	1000	1000	1000
3	1000	1400	1400	1000	1200	1400
4	1200	1400	1600	1200	1400	1400
5	1200	1400	1600	1200	1400	1600
6	1400	1600	1800	1200	1400	1600
7	1400	1600	1800	1200	1600	1800
8	1400	1600	2000	1400	1600	1800
9	1600	1800	2000	1400	1600	1800
10	1600	1800	2200	1400	1800	2000
11	1800	2000	2200	1600	1800	2000
12	1800	2200	2400	1600	2000	2200
13	2000	2200	2600	1600	2000	2200
14	2000	2400	2800	1800	2000	2400
15	2200	2600	3000	1800	2000	2400
16	2400	2800	3200	1800	2000	2400
17	2400	2800	3200	1800	2000	2400
18	2400	2800	3200	1800	2000	2400
19-20	2600	2800	3000	2000	2200	2400
21-25	2400	2800	3000	2000	2200	2400
26-30	2400	2600	3000	1800	2000	2400
31-35	2400	2600	3000	1800	2000	2200
36-40	2400	2600	2800	1800	2000	2200
41-45	2200	2600	2800	1800	2000	2200
46-50	2200	2400	2800	1800	2000	2200
51-55	2200	2400	2800	1600	1800	2200
56-60	2200	2400	2600	1600	1800	2200
61-65	2000	2400	2600	1600	1800	2000
66-70	2000	2200	2600	1600	1800	2000
71-75	2000	2200	2600	1600	1800	2000
76 and up	2000	2200	2400	1600	1800	2000

¹Based on Estimated Energy Requirements (EER) equations, using reference heights (average) and reference weights (healthy) for each age/sex group, rounded to the nearest 200 calories. EER equations are from the Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. Washington DC: National Academies Press, 2002.

Source: Britten et al., 2006.

Table B2.2. Top five sources of energy among US children, adolescents, and adults by age, NHANES 2005-06¹

	Overall, ages 2+ years	Ages 2-18 years	Ages 2-3 years	Ages 4-8 years	Ages 9-13 years	Ages 14-18 years	Ages 19+ years
Mean energy intake (kcal)	2157	2027	1471	1802	2035	2427	2199
Rank							
1	Grain-based desserts ¹ (138 kcal)	Grain-based desserts (138 kcal)	Whole milk (104 kcal)	Grain-based desserts (136 kcal)	Grain-based desserts (145 kcal)	Soda/energy/sports drinks ² (226 kcal)	Grain-based desserts (138 kcal)
2	Yeast breads (129 kcal)	Pizza (136 kcal)	100% fruit juice (not orange or grapefruit) (93 kcal)	Yeast breads (98 kcal)	Pizza (128 kcal)	Pizza (213 kcal)	Yeast breads (134 kcal)
3	Chicken and chicken mixed dishes (121 kcal)	Soda/energy/sports drinks (118 kcal)	Reduced fat milk (91 kcal)	Pasta and pasta dishes (97 kcal)	Chicken and chicken mixed dishes (122 kcal)	Grain-based desserts (157 kcal)	Chicken and chicken mixed dishes (123 kcal)
4	Soda/energy/sports drinks (114 kcal)	Yeast breads (114 kcal)	Pasta and pasta dishes (86 kcal)	Pizza (95 kcal)	Yeast breads (109 kcal)	Yeast breads (151 kcal)	Soda/energy/sports drinks ² (112 kcal)
5	Pizza (98 kcal)	Chicken and chicken mixed dishes (113 kcal)	Grain-based desserts (68 kcal)	Reduced fat milk (95 kcal)	Soda/energy/sports drinks (105 kcal)	Chicken and chicken mixed dishes (143 kcal)	Alcoholic beverages (106 kcal)

¹Foods ranked by mean contribution to overall energy intake. Table shows each food category and its mean caloric contribution for each age group.

²Includes cakes, cookies, doughnuts, pies, crisps, cobblers, granola bars.

³Includes sodas, energy drinks, sports drinks, and sweetened bottled water including vitamin water.

Note: For a more detailed listing of food sources of energy, see **Part D. Section 1. Energy Balance**, Tables D1.1, D1.6, and D1.7.

Source: National Cancer Institute (NCI). Food Sources of Energy Among US Population, 2005-06. Risk Factor Monitoring and Methods Branch Website. Applied Research Program. National Cancer Institute, 2010a. <http://riskfactor.cancer.gov/diet/foodsources/>. Updated May 21, 2010. Accessed May 21, 2010.

Table B2.3. USDA Food Patterns—recommended daily intake amounts¹ from each food group or subgroup at all calorie levels. Recommended intakes from vegetable subgroups are per week

Energy level of pattern ²	1000	1200	1400	1600	1800	2000	2200	2400	2600	2800	3000	3200
Fruits	1 c	1 c	1½ c	1½ c	1½ c	2 c	2 c	2 c	2 c	2½ c	2½ c	2½ c
Vegetables	1 c	1½ c	1½ c	2 c	2½ c	2½ c	3 c	3 c	3½ c	3½ c	4 c	4 c
Dark green vegetables	½ c/wk	1 c/wk	1 c/wk	1 ½ c/wk	1 ½ c/wk	1 ½ c/wk	2 c/wk	2 c/wk	2 ½ c/wk	2 ½ c/wk	2 ½ c/wk	2 ½ c/wk
Red/Orange vegetables	2½ c/wk	3 c/wk	3 c/wk	4 c/wk	5 ½ c/wk	5 ½ c/wk	6 c/wk	6 c/wk	7 c/wk	7 c/wk	7½ c/wk	7½ c/wk
Cooked dry beans and peas	½ c/wk	½ c/wk	½ c/wk	1 c/wk	1 ½ c/wk	1 ½ c/wk	2 c/wk	2 c/wk	2 ½ c/wk	2 ½ c/wk	3 c/wk	3 c/wk
Starchy vegetables	2 c/wk	3½ c/wk	3½ c/wk	4 c/wk	5 c/wk	5 c/wk	6 c/wk	6 c/wk	7 c/wk	7 c/wk	8 c/wk	8 c/wk
Other vegetables	1½ c/wk	2½ c/wk	2½ c/wk	3½ c/wk	4 c/wk	4 c/wk	5 c/wk	5 c/wk	5½ c/wk	5½ c/wk	7 c/wk	7 c/wk
Grains	3 oz eq	4 oz eq	5 oz eq	5 oz eq	6 oz eq	6 oz eq	7 oz eq	8 oz eq	9 oz eq	10 oz eq	10 oz eq	10 oz eq
Whole grains	1½ oz eq	2 oz eq	2½ oz eq	3 oz eq	3 oz eq	3 oz eq	3½ oz eq	4 oz eq	4½ oz eq	5 oz eq	5 oz eq	5 oz eq
Other grains	1½ oz eq	2 oz eq	2½ oz eq	2 oz eq	3 oz eq	3 oz eq	3½ oz eq	4 oz eq	4½ oz eq	5 oz eq	5 oz eq	5 oz eq
Meat and beans	2 oz eq	3 oz eq	4 oz eq	5 oz eq	5 oz eq	5½ oz eq	6 oz eq	6 ½ oz eq	6 ½ oz eq	7 oz eq	7 oz eq	7 oz eq
Milk	2 c	2 c	2 c	3 c	3 c	3 c	3 c	3 c	3 c	3 c	3 c	3 c
Oils	15 g	17 g	17 g	22 g	24 g	27 g	29 g	31 g	34 g	36 g	44 g	51g
Maximum SoFAS ³ limit, calories (%total calories)	137 (14%)	137 (11%)	137 (10%)	121(8%)	161(9%)	258 (13%)	266 (12%)	330 (14%)	362 (14%)	395 (14%)	459 (15%)	596 (19%)

¹Food group amounts shown in cup (c) or ounce equivalents (oz eq). Oils are shown in grams (g). Quantity equivalents for each food group are:

- Grains, 1 ounce equivalent is: ½ cup cooked rice, pasta, or cooked cereal; 1 ounce dry pasta or rice; 1 slice bread; 1 small muffin (1 oz); 1 oz ready-to-eat cereal.
- Fruits and vegetables, 1 cup equivalent is: 1 cup raw or cooked fruit or vegetable, 1 cup fruit or vegetable juice, 2 cups leafy salad greens.
- Meat and beans, 1 ounce equivalent is: 1 ounce lean meat, poultry, fish; 1 egg; ¼ cup cooked dry beans; 1 Tbsp peanut butter; ½ ounce nuts/ seeds.
- Milk, 1 cup equivalent is: 1 cup milk or yogurt, 1½ ounces natural cheese such as Cheddar cheese or 2 ounces of processed cheese.

²Food intake patterns at 1000, 1200, and 1400 calories meet the nutritional needs of children ages 2 to 8 years. Patterns from 1600 to 3200 calories meet the nutritional needs of children 9 years of age and older and adults. If a child ages 2 to 8 years needs more calories and, therefore, is following a pattern at 1600 calories or more, the recommended amount from the milk group can be 2 cups per day. Children ages 9 years and older and adults should not use the 1000, 1200, or 1400 calorie patterns.

³SoFAS are calories from solid fats and added sugars.

Table B2.4. Dietary pattern comparison: Current US Intake, DASH-sodium diet, and USDA food patterns. Description, nutrient composition, and food group amounts (adjusted to 2000 calories)

Dietary Pattern	Usual US Intake Adults	DASH with Reduced Sodium	USDA Base Pattern ¹	USDA Plant-based	USDA Lacto-ovo Vegetarian	USDA Vegan
Citation	NHANES 2001-04; 2005-06; Ages 19+	Karanja et al., 1999 and Lin et al., 2003	Britten et al., 2006; Online Appendix E-3.1	Online Appendix E-3.3	Online Appendix E-3.3	Online Appendix E-3.3
Qualitative Description						
Emphasizes		Potassium-rich vegetables, fruits, and low-fat milk products	Vegetables, fruits, and whole grains, low-fat milk products	Plant foods - vegetables, fruits, whole grains, legumes, low-fat milk products	Plant foods - vegetables, fruits, whole grains, legumes, nuts, seeds, soy foods, milk products	Plant foods - vegetables, fruits, whole grains, legumes, nuts, seeds, soy foods
Includes		Whole grains, poultry, fish, and nuts	Enriched grains, lean meat, fish, and oils	Lean meat, eggs, fish, and oils	Eggs, oils	Non-dairy milk alternatives
Limits (small amount)		Red meats, sweets, and sugar-containing beverages	Solid fats Added sugars	Solid fats Added sugars	No meat, poultry, fish Added sugars	No animal products Added sugars
Nutrients						
Calories (kcal)	2000	2000	2000	2000	2000	2000
Carbohydrates (% total kcal)	48.4%	58%	56.7%	55.8%	56.7%	56.8%
Protein (% total kcal)	15.2%	18%	15.2%	16.3%	15.2%	13.3%
Total Fat (% total kcal)	33.5%	27%	32%	31%	31%	33%
Saturated Fat (% total kcal)	10.9%	6%	8.4%	7.8%	7.8%	6.8%
Monounsaturated (% total kcal)	12.5%	10%	12.0%	11.4%	11.8%	12.4%
Polyunsaturated (% total kcal)	6.8%	8%	9.0%	9.3%	9.4%	12.0%
Cholesterol (mg)	269	143	229	170	160	17
Fiber (g)	15	29	30	37	39	43
Potassium (mg)	2909	4371	3478	3611	3610	3645
Sodium (mg)	2846	1095	1722	1582	1595	1224
Food Groups						
Vegetables: total (c)	1.6	2.1	2.5	2.5	2.5	2.5
- Dark Green (c)	0.1	nd	0.2	0.2	0.2	0.2
- Legumes ² (c)	0.1	nd	0.2	0.2	0.2	0.2
- Red Orange (c)	0.4	nd	0.8	0.8	0.8	0.8
- Other Veg (c)	0.5	nd	0.6	0.6	0.6	0.6
- Starchy Veg (c)	0.5	nd	0.7	0.7	0.7	0.7

Table B2.4 (continued). Dietary pattern comparison: Current US Intake, DASH-sodium diet, and USDA food patterns. Description, nutrient composition, and food group amounts (adjusted to 2000 calories)

Dietary Pattern	Usual US Intake Adults 19 year+	DASH with Reduced Sodium	USDA Base Pattern ¹	USDA Plant-based	USDA Lacto-ovo Vegetarian	USDA Vegan
Calories (kcal)	2000	2000	2000	2000	2000	2000
Food Groups						
Fruit & juices (c)	1.0	2.5	2	2	2	2
Grains: total (oz)	6.4	7.3	6	6	6	6
- Whole grains (oz)	0.6	3.9	3	3	3	3
Milk & milk products incl whole fat (c)	1.5	0.7 (whole)	-	-	-	-
- Low-fat milk (c)	Nd	1.9	3	3	3	3 (non-dairy) ³
Animal Proteins:						
- Meat (oz)	2.5	1.4	2.5	0.6	-	-
- Poultry (oz)	1.2	1.7	1.5	0.4	-	-
- Eggs (oz)	0.4	nd	0.4	0.4	0.6	-
- Fish (total) (oz)	0.5	1.4	0.5	0.7	-	-
-- Hi N3 (oz)	0.1	nd	0.1	nd	-	-
-- Low N3 (oz)	0.4	nd	0.4	nd	-	-
Plant Proteins:						
- Legumes (oz)	nd	0.4	See vegetables.	1.4	1.4	1.9
- Nuts & seeds (oz)	0.5	0.9	0.6	1.1	1.9	2.1
- Soy products (oz)	0.0	nd	0.05	0.9	1.7	1.4
Oils (g)	17.7	24.8	27	23	19	18
Solid Fats (g)	43.2	nd	16	16	16	16
Added Sugar (g)	79.0	12 (snacks/sweets)	32	32	32	32
Alcohol (g)	9.9	-	-	-	-	-

¹ The USDA Base Food Pattern is slightly adapted from the 2000-calorie pattern presented in the 2005 Dietary Guidelines for Americans (DGA). Vegetable subgroups were realigned to include a Red/Orange subgroup. The base pattern and the vegetarian variations are subject to change as the 2010 DGA are developed. The measures are cup and ounce equivalents (Britten, 2006; Marcoe, 2006). Nutrient distribution updated with 2010 composites.

² On USDA patterns, total recommended legume amount is the sum of amounts recommended in the Vegetable and the Meat & Beans groups. An ounce equivalent of legumes in the Meat & Beans group is ¼ cup. For example, in the 2000-calorie pattern, total weekly legume recommendation is (13 oz eq /4) + 1.5 cups = 5 cups.

³ Non-dairy options in Vegan pattern are calcium-fortified soymilk, rice milk, and tofu. All USDA patterns contain a small amount of soy milk.

nd = Not described.

(-) = No recommendation.

Sources: Usual US Intakes – WWEIA, NHANES 2001-2004 and WWEIA, NHANES 2005-2006, one-day mean intakes consumed per individual. Male and female intakes adjusted to 2000 calories, averaged, and rounded to one decimal point.

Part B. Section 2. Appendix: Dietary Patterns and Health Outcomes

Introduction

Across the world and within the United States, there are striking differences in diet. Concomitantly, there are substantial differences in health outcomes, many of which are related to diet. This section discusses several dietary patterns that are associated with desirable health outcomes. It focuses on total mortality, cardiovascular disease (CVD), and blood pressure, a major diet-related cardiovascular risk factor. The World Cancer Research Fund/American Institute for Cancer Research, recently reviewed the available evidence of the relationship of cancer with specific dietary factors and overall dietary patterns (WCRF/AICR, 2007). Although several dietary factors were associated with specific types of cancer, it concluded that no firm judgment can be made on the relationship of dietary patterns with cancer, in large part, because variability in definitions precluded a formal synthesis of evidence.

The study of dietary patterns is complex. First, there is substantial heterogeneity even among diets that fall under a common rubric (e.g., Mediterranean diets). Second, dietary patterns are not static. Traditional diets known for their health benefits (e.g., Mediterranean and Okinawan diets) are being supplanted by versions that often reflect Western culture and that lead to worse not better health outcomes. For this reason, we focused on pre-transition dietary patterns. Third, with few exceptions, standardized assessment of diet is unavailable, making it difficult to compare diets. Fourth, health outcomes are often unavailable and, when available, are not directly comparable across studies. Fifth, dietary patterns, even with proven health benefits, may not be ideal and could be improved. For example, traditional Japanese diets are associated with a low risk of coronary heart disease but a high risk of stroke, likely because of excessive sodium intake. Sixth, describing dietary patterns and evaluating their health outcomes often requires scoring systems based on adherence to specific aspects of the diets. This approach commonly relies on researchers who exercise best judgment in selecting biologically relevant aspects of the diet and in developing a formula, which typically weights each dimension as of equivalent importance. Seventh, in the interpretation of observational data, particularly ecologic data, it is difficult to separate the effects of diet from other factors, such as smoking and physical inactivity, that likely account for part of the observed differences in health outcomes.

Despite these caveats, the 2010 Dietary Guidelines Advisory Committee (DGAC) was able to identify dietary patterns that are associated with substantial beneficial health benefits (Table B2.5). Specifically, the Committee focused on the following dietary patterns for which there was research on health outcomes as well as information on nutrient and food group composition: 1) Dietary

Approaches to Stop Hypertension (DASH)-style dietary patterns, 2) Mediterranean-style dietary patterns, and 3) Vegetarian dietary patterns. The DASH dietary pattern is a Western-style dietary pattern for which a large and burgeoning literature documents its health benefits. The Committee also included Mediterranean and Japanese dietary patterns, which were associated with the lowest risk of coronary heart disease in the Seven Countries study (Keys, 1980). Subsequently, a substantial literature has documented the health benefits of Mediterranean-style diets. In contrast, while traditional Asian dietary patterns (e.g., Japanese and Okinawan dietary patterns) have also been associated with a reduced risk of coronary heart disease (Wilcox, 2007), documentation using contemporary research methods is scant. Finally, the Committee studied vegetarian diets, which have been associated with a reduced risk of coronary heart disease (Key, 1999).

DASH-style Dietary Patterns

DASH-style dietary patterns emphasize fruits, vegetables, and low-fat dairy products; include whole grains, poultry, fish and nuts; and are reduced in red meat, sweets, and sugar-containing beverages (Karanja, 1999; Craddick, 2003). The diets are rich in potassium, magnesium, calcium and fiber, and reduced in saturated fat and cholesterol. As originally tested, the DASH diet is reduced in total fat (27% kcal) with total protein intake of 18 percent of calories and carbohydrate intake of 55 percent of calories. However, other versions of the DASH diet are available, in which carbohydrate is partially replaced with protein (about half from plant sources) or unsaturated fat (predominantly monounsaturated fat) (Appel, 2005; Swain, 2008). The latter version is noteworthy because nutrient adequacy and a reduced saturated fat intake (6% kcal) were both achieved in the setting of high monounsaturated fat intake (21% kcal). Each of these DASH-style diets lowers blood pressure, improves blood lipids, and reduces CVD risk. Blood pressure reduction is the greatest when the DASH diet is consumed with reduced sodium intake (Sacks, 2001).

As originally developed, the DASH diet was designed to provide a nutrient profile that might lower blood pressure. As such, it is a derived dietary pattern. Nonetheless, it is based on foods that are routinely available in US and was studied using foods purchased at local stores. At present, few adults, even those with hypertension, eat a diet that is consistent with the DASH dietary pattern (Mellen, 2008).

Mediterranean-style Dietary Patterns

In view of the large number of cultures and agricultural patterns of countries that border the Mediterranean Sea, the “Mediterranean” diet is not a single dietary pattern. Countries included those

of southern-most Europe, the Middle East, and northern-most Africa. Interest in traditional Mediterranean-style diets is substantial because such diets have been associated with considerable health benefits. Because of the multiplicity of dietary patterns termed “Mediterranean,” it has been challenging to characterize these diets. Although a traditional Mediterranean diet has no well-accepted set of criteria, it can be described as one that emphasizes breads and other cereal foods usually made from wheat, vegetables, fruits, nuts, unrefined cereals, and olive oil; includes fish and wine with meals (in non-Islamic countries); and is reduced in saturated fat, meat, and full-fat dairy products (Kris-Etherton, 2001; Trichopoulou, 2003; WCRF/AICR, 2007). Table B2.5 displays the nutrient profile and food group composition of Mediterranean-style diets, as reported in three cohort studies (one from Greece, one from Spain, and one from the US) (Fung, 2009; Karanja, 1999; Lin, 2003; Nunez-Cordoba, 2008; Trichopoulou, 2003; Wilcox, 2007).

Results from observational studies and clinical trials suggest that consumption of a traditional Mediterranean diet, similar to that of Crete in the 1960s, is associated with one of the lowest risks of coronary heart disease in the world. Over time, the diet of Crete has changed remarkably and is now characterized by higher intake of saturated fat and cholesterol, and reduced intake of monounsaturated fats. At the same time, total fat consumption has fallen. These trends have been accompanied by a steady rise in coronary heart disease risk (Menotti, 1999).

Vegetarian Dietary Patterns

In many observational studies, vegetarian diets and lifestyle have been associated with improved health outcomes. The types of vegetarian diets consumed in the US vary considerably. Strict vegetarians (i.e., vegans), do not consume any animal products, while other types of vegetarians, such as lacto-ovo vegetarians, consume milk and eggs. Although not strict vegetarians, many individuals consume small or minimal amounts of animal products. On average, vegetarians consume fewer calories from fat than non-vegetarians, particularly saturated fat, and have a higher consumption of carbohydrates than non-vegetarians. In addition, vegetarians tend to consume fewer overall calories and have a lower body mass index than non-vegetarians. These characteristics, in addition to the dietary pattern per se, may contribute to the improved health outcomes of vegetarians.

Although no or minimal consumption of animal products is a hallmark of vegetarian diets, these diets have a clear potential for confounding, particularly from other dietary and non-dietary factors. Hence, the improved health experience of vegetarians may not only result from reduced consumption of saturated fats but also from greater consumption of vegetables, fruit, nuts, and grains or from other health attributes, such as not smoking cigarettes).

Other Dietary Patterns

In view of the increasing diversity of the US population, interest in the health effects of non-Western diets is substantial. One group of diets with potential health benefits are those consumed in Asia. It is well-documented that in Southeast Asia, coronary heart disease rates have been among the lowest in the world. Lifestyle factors, especially diet, appear to be a major reason. However, contemporary evidence (e.g., prospective cohort studies and clinical trials), similar to the evidence available for the other types of diets is sparse.

Traditional Japanese dietary patterns emphasize soybean products, fish, seaweeds, vegetables, fruit, and green tea, and are reduced in meats (Shimazu, 2007). Nonetheless, it should be recognized that this diet is high in salt, likely accounting for the high incidence of stroke in this population. Similar to other dietary patterns, Japanese dietary patterns have evolved over time.

The longevity of Okinawans is among the highest in the world. Researchers attribute the longevity and health of Okinawans, in large part, to diet composition or some other aspect of their diet, such as energy restriction (Willcox, 2007). The indigenous Satsamu sweet potato, which is rich in nutrients, is the food staple that provides the bulk of energy intake. Other prominent foods are a wide variety of seaweeds, Okinawan tofu, and herbaceous plants. Okinawan food culture also includes a modest amount of fish and pork. The estimated carbohydrate content of this diet is extremely high, at more than 80 percent of calories. Salt intake is the lowest of all Japan. However, the traditional Okinawan diet has changed such that fast foods and processed foods are increasingly consumed.

What is the Effect of Different Dietary Patterns (DASH, Mediterranean, Vegetarian, and Other) on Blood Pressure in Adults?

The 2010 DGAC performed a literature search to identify research, with no date limits, on the effect of the above dietary patterns on blood pressure in adults. Some articles were reviewed that included dietary patterns that were characterized using dietary cluster or factor analysis. The NEL search identified 146 potential articles (11 reviews/meta-analyses and 135 primary studies). Of these, 126 were excluded. A total of 20 articles, all of them primary studies, met the eligibility criteria and were reviewed (Table B2.6).

Of the 12 studies that evaluated a DASH-style dietary pattern (Appel, 2005; Appel, 1997; Appel, 2003; Azadbakht, 2005; Dauchet, 2007; Forman, 2009; Miller, 2002; Nowson, 2009; Nowson, 2005; Nowson, 2004; Sacks, 2001; Schulze, 2003), nine were randomized controlled trials (Appel, 2005;

Appel, 1997; Appel, 2003; Azadbakht, 2005; Miller, 2002; Nowson, 2009; Nowson, 2005; Nowson, 2004; Sacks, 2001), and three were prospective cohort studies (Dauchet, 2007; Forman, 2009; Schulze, 2003). In aggregate, the DASH diet lowered systolic blood pressure in 12 studies (Appel, 2005; Appel, 1997; Appel, 2003; Azadbakht, 2005; Dauchet, 2007; Forman, 2009; Miller, 2002; Nowson, 2009; Nowson, 2005; Nowson, 2004; Sacks, 2001; Schulze, 2003) and diastolic blood pressure in 10 of the 12 studies that reported diastolic blood pressure (Appel, 2005; Appel, 1997; Appel, 2003; Azadbakht, 2005; Dauchet, 2007; Forman, 2009; Miller, 2002; Nowson, 2005; Nowson, 2004; Schulze, 2003). In several instances, blood pressure reduction occurred as part of a multi-factorial intervention that tested the DASH dietary pattern concomitantly with other interventions (Appel, 2003; Miller, 2002; Sacks, 2001).

Few studies examined the effects of a Mediterranean-style diet on blood pressure. In the one available study (Núñez-Córdoba, 2009) a cohort study, a Mediterranean-style diet, lowered systolic and diastolic blood pressure.

Four trials tested the effects of vegetarian diets on blood pressure (Hakala and Karvetti, 1989; Margetts, 1986; Rouse, 1983; Sciarrone 1993). Vegetarian-style dietary patterns lowered systolic blood pressure in all four trials and diastolic blood pressure in three trials (Hakala and Karvetti, 1989; Rouse, 1983; Sciarrone, 1993).

One randomized, cross-over trial found that, within the context of a traditional Japanese diet, increased vegetables and fruit intake and decreased sodium intake significantly reduced systolic blood pressure in normotensive and hypertensive free-living rural Japanese (Takahashi, 2006).

What is the Effect of Different Dietary Patterns (DASH, Mediterranean, Vegetarian, and Other) on Cardiovascular Disease, Stroke, and Total Mortality in Adults?

The 2010 DGAC performed a literature search to identify research, with no date limits, on the effect of these dietary patterns on cardiovascular disease, stroke, and total mortality in adults. Some articles were reviewed that included dietary patterns that were characterized using dietary clusters or factor analysis. The search identified 197 potential articles (11 reviews/meta-analyses and 186 primary studies). Of these, 168 were excluded. A total of 29 articles, 27 primary studies, one systematic review/meta-analysis, and one systematic review, met the eligibility criteria and were reviewed. Of the 27 primary studies, two were randomized controlled trials, 20 were prospective cohort studies (two were follow-up of RCTs and one was non-concurrent), three were case-control studies, one was a med adherence analysis, and one was a time series (Table B2.7).

Of the 10 studies that evaluated a DASH-style dietary pattern, nine were prospective cohort studies (Folsom, 2007; Fung, 2001; Fung, 2008; Heidemann, 2008; Hu, 2000; Levitan, 2009; Osler, 2001; Parikh, 2009; Singman, 1980) and one was a randomized trial in which estimated coronary heart disease risk was the outcome (Appel, 2005). Of the 10 that evaluated a relationship of a DASH-style dietary pattern with CVD, nine studies documented that consumption of a DASH-style diet was associated with a reduced risk of CVD (Appel, 2005; Fung, 2001; Fung, 2008; Heidemann, 2008; Hu, 2000; Levitan, 2009; Osler, 2001; Parikh, 2009; Singman, 1980), and one (Folsom, 2007) found no such relationship. For total mortality, six of seven studies that reported data on mortality documented an inverse relation (Fung, 2008; Heidemann, 2008; Hu, 2000; Levitan, 2009; Osler, 2001; Parikh, 2009) and one (Folsom, 2007) found no such relationship. In the two available studies with stroke (Fung, 2008; Parikh, 2009), consumption of a DASH-style pattern prevented stroke.

Several studies examined the effects of a Mediterranean style diet on CVD and total mortality. Of the 13 studies, one was a systematic review/meta-analysis (Mente, 2009), one was a meta-analysis (Sofi, 2008), nine were prospective cohort studies (Fidanza, 2004; Fung, 2009; Harriss, 2007; Knoops, 2004; Mitrou, 2007; Panagiotakos, 2009; Trichopoulou, 2003; Trichopoulou, 2009; Waijers, 2006), one was an adherence analysis (Alberti, 2008), and one was a case-control study (Panagiotakos, 2005). Of the 10 studies that evaluated a relationship of a Mediterranean-style dietary pattern with CVD, each documented a beneficial effect (Fidanza, 2004; Fung, 2009; Harriss, 2007; Knoops, 2004; Mente, 2009; Mitrou, 2007; Panagiotakos, 2009; Panagiotakos, 2005; Sofi, 2008; Trichopoulou, 2003). Likewise, of the 10 studies with data on total mortality, each documented an inverse relation (Alberti, 2008; Fidanza, 2004; Fung, 2009; Harriss, 2007; Knoops, 2004; Mitrou, 2007; Sofi, 2008; Trichopoulou, 2003; Trichopoulou, 2009; Waijers, 2006). In the one available study with stroke, consumption of a Mediterranean-style pattern prevented stroke (Fung, 2009).

Five studies examined the effects of a vegetarian diet on CVD and total mortality. Of the five studies, three were prospective cohort studies (Chang-Claude, 2005; Key, 1996; Mann, 1997), one was a meta-analysis (Key, 1998), and one was a time series analysis (Fraser, 2005). Of the five studies with CVD as the study outcome, all found that vegetarian diets were associated with a reduced risk of CVD compared to non-vegetarian diets (Chang-Claude, 2005; Fraser, 2005; Key, 1998; Key, 1996; Mann, 1997). For total mortality, four studies (Fraser, 2005; Key, 1998; Key, 1996; Mann, 1997) documented that a vegetarian diet was associated with a reduced risk of death, and one study (Chang-Claude, 2005) did not detect an association.

One prospective cohort study (Shimazu, 2007) assessed the association between dietary patterns among the Japanese and CVD mortality. Three diet patterns were identified: 1) Japanese pattern including soybean products, fish, seaweed, vegetables, fruit and green tea, 2) animal food pattern, and 3) high-dairy, high-fruit and vegetable, low alcohol (DFA) pattern. The Japanese pattern was

associated with a decreased risk of CVD mortality, while the animal food pattern was associated with increased risk. The DFA pattern was not significantly associated with a change in CVD risk.

Conclusion

The totality of evidence documenting a beneficial impact of plant-based, lower-sodium dietary patterns on CVD risk is remarkable. Indeed, several distinct dietary patterns are associated with lower blood pressure and a reduced risk of CVD and total mortality. When explicitly tested, a reduced sodium intake further lowers blood pressure. A common feature of these diets is an emphasis on plant-based foods. Accordingly, fiber intake is high while saturated fat typically low. When total fat intake is high, that is, over 30 percent of calories, the predominant fat is monounsaturated or polyunsaturated fat. Carbohydrate intake is often, but not necessarily high; the predominant forms appear to be complex carbohydrates, often from whole grain products with minimal processing.

References

Alberti A, Fruttini D, Fidanza F. The Mediterranean Adequacy Index: further confirming results of validity. *Nutr Metab Cardiovasc Dis*. 2009 Jan;19(1):61-6.

Appel LJ, Sacks FM, Carey VJ, Obarzanek E, Swain JF, Miller ER 3rd, Conlin PR, Erlinger TP, Rosner BA, Laranjo NM, Charleston J, McCarron P, Bishop LM; OmniHeart Collaborative Research Group. Effects of protein, monounsaturated fat, and carbohydrate intake on blood pressure and serum lipids: results of the OmniHeart randomized trial. *JAMA*. 2005 Nov 16;294(19):2455-64.

Appel LJ, Champagne CM, Harsha DW, Cooper LS, Obarzanek E, Elmer PJ, Stevens VJ, Vollmer WM, Lin PH, Svetkey LP, Stedman SW, Young DR; Writing Group of the PREMIER Collaborative Research Group. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *JAMA*. 2003 Apr 23-30;289(16):2083-93.

Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, Bray GA, Vogt TM, Cutler JA, Windhauser MM, Lin PH, Karanja N. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med*. 1997 Apr 17;336(16):1117-24.

Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi T, Azizi F. Beneficial effects of a Dietary Approaches to Stop Hypertension eating plan on features of the metabolic syndrome. *Diabetes Care*. 2005 Dec;28(12):2823-31.

Brunner EJ, Mosdøl A, Witte DR, Martikainen P, Stafford M, Shipley MJ, Marmot MG. Dietary patterns and 15-y risks of major coronary events, diabetes, and mortality. *Am J Clin Nutr*. 2008 May;87(5):1414-21.

Cai H, Shu XO, Gao YT, Li H, Yang G, Zheng W. A prospective study of dietary patterns and mortality in Chinese women. *Epidemiology*. 2007 May;18(3):393-401.

Chang-Claude J, Hermann S, Eilber U, Steindorf K. Lifestyle determinants and mortality in German vegetarians and health-conscious persons: results of a 21-year follow-up. *Cancer Epidemiol Biomarkers Prev*. 2005 Apr;14(4):963-8.

Craddick SR, Elmer PJ, Obarzanek E, Vollmer WM, Svetkey LP, Swain MC. The DASH diet and blood pressure. *Curr Atheroscler Rep*. 2003 Nov;5(6):484-91.

Dauchet L, Kesse-Guyot E, Czernichow S, Bertrais S, Estaquio C, Péneau S, Vergnaud AC, Chat-Yung S, Castetbon K, Deschamps V, Brindel P, Hercberg S. Dietary patterns and blood pressure change over 5-y follow-up in the SU.VI.MAX cohort. *Am J Clin Nutr*. 2007 Jun;85(6):1650-6.

Fidanza F, Alberti A, Lanti M, Menotti A. Mediterranean Adequacy Index: correlation with 25-year mortality from coronary heart disease in the Seven Countries Study. *Nutr Metab Cardiovasc Dis*. 2004 Oct;14(5):254-8.

Folsom AR, Parker ED, Harnack LJ. Degree of concordance with DASH diet guidelines and incidence of hypertension and fatal cardiovascular disease. *Am J Hypertens*. 2007 Mar;20(3):225-32.

Forman JP, Stampfer MJ, Curhan GC. Diet and lifestyle risk factors associated with incident hypertension in women. *JAMA*. 2009 Jul 22;302(4):401-11.

Fraser GE. A comparison of first event coronary heart disease rates in two contrasting California populations. *J Nutr Health Aging*. 2005;9(1):53-8.

Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G, Hu FB. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med*. 2008 Apr 14;168(7):713-20. Erratum in: *Arch Intern Med*. 2008 Jun 23;168(12):1276.

Fung TT, Rexrode KM, Mantzoros CS, Manson JE, Willett WC, Hu FB. Mediterranean diet and incidence of and mortality from coronary heart disease and stroke in women. *Circulation*. 2009 Mar 3;119(8):1093-100.

Fung TT, Willett WC, Stampfer MJ, Manson JE, Hu FB. Dietary patterns and the risk of coronary heart disease in women. *Arch Intern Med*. 2001 Aug 13-27;161(15):1857-62.

Hakala P, Karvetti RL. Weight reduction on lactovegetarian and mixed diets. Changes in weight, nutrient intake, skinfold thicknesses and blood pressure. *Eur J Clin Nutr*. 1989 Jun;43(6):421-30.

Harriss LR, English DR, Powles J, Giles GG, Tonkin AM, Hodge AM, Brazionis L, O'Dea K. Dietary patterns and cardiovascular mortality in the Melbourne Collaborative Cohort Study. *Am J Clin Nutr*. 2007 Jul;86(1):221-9.

Heidemann C, Schulze MB, Franco OH, van Dam RM, Mantzoros CS, Hu FB. Dietary patterns and risk of mortality from cardiovascular disease, cancer, and all causes in a prospective cohort of women. *Circulation*. 2008 Jul 15;118(3):230-7.

Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC. Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr*. 2000 Oct;72(4):912-21.

Karanja NM, Obarzanek E, Lin PH, McCullough ML, Phillips KM, Swain JF, Champagne CM, Hoben KP. Descriptive characteristics of the dietary patterns used in the Dietary Approaches to Stop Hypertension Trial. DASH Collaborative Research Group. *J Am Diet Assoc*. 1999 Aug;99(8 Suppl):S19-27.

Key TJ, Fraser GE, Thorogood M, Appleby PN, Beral V, Reeves G et al. Mortality in vegetarians and nonvegetarians: detailed findings from a collaborative analysis of 5 prospective studies. *Am J Clin Nutr*. 1999; 70(3 Suppl):516S-524S.

Key TJ, Fraser GE, Thorogood M, Appleby PN, Beral V, Reeves G, Burr ML, Chang-Claude J, Frentzel-Beyme R, Kuzma JW, Mann J, McPherson K. Mortality in vegetarians and non-vegetarians: a collaborative analysis of 8300 deaths among 76,000 men and women in five prospective studies. *Public Health Nutr*. 1998 Mar;1(1):33-41.

- Key TJ, Thorogood M, Appleby PN, Burr ML. Dietary habits and mortality in 11,000 vegetarians and health conscious people: results of a 17 year follow up. *BMJ*. 1996 Sep 28;313(7060):775-9.
- Keys A. *Seven countries: a multivariate analysis of death and coronary heart disease*. Cambridge, MA.: Harvard University Press, 1980.
- Knoops KT, de Groot LC, Kromhout D, Perrin AE, Moreiras-Varela O, Menotti A, van Staveren WA. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA*. 2004 Sep 22;292(12):1433-9.
- Kris-Etherton P, Eckel RH, Howard BV, St Jeor S, Bazzarre TL; Nutrition Committee Population Science Committee and Clinical Science Committee of the American Heart Association. AHA Science Advisory: Lyon Diet Heart Study. Benefits of a Mediterranean-style, National Cholesterol Education Program/American Heart Association Step I Dietary Pattern on Cardiovascular Disease. *Circulation*. 2001 Apr 3;103(13):1823-5.
- Levitan EB, Wolk A, Mittleman MA. Consistency with the DASH diet and incidence of heart failure. *Arch Intern Med*. 2009 May 11;169(9):851-7.
- Mann JI, Appleby PN, Key TJ, Thorogood M. Dietary determinants of ischaemic heart disease in health conscious individuals. *Heart*. 1997 Nov;78(5):450-5.
- Margetts BM, Beilin LJ, Vandongen R, Armstrong BK. Vegetarian diet in mild hypertension: a randomised controlled trial. *Br Med J (Clin Res Ed)*. 1986 Dec 6;293(6560):1468-71.
- Mellen PB, Gao SK, Vitolins MZ, Goff DC Jr. Deteriorating dietary habits among adults with hypertension: DASH dietary accordance, NHANES 1988-1994 and 1999-2004. *Arch Intern Med*. 2008 Feb 11;168(3):308-14.
- Mente A, de Koning L, Shannon HS, Anand SS. A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med*. 2009 Apr 13;169(7):659-69.
- Menotti A, Kromhout D, Blackburn H, Fidanza F, Buzina R, Nissinen A. Food intake patterns and 25-year mortality from coronary heart disease: cross-cultural correlations in the Seven Countries Study. The Seven Countries Study Research Group. *Eur J Epidemiol*. 1999; 15(6):507-515.
- Miller ER 3rd, Erlinger TP, Young DR, Jehn M, Charleston J, Rhodes D, Wasan SK, Appel LJ. Results of the Diet, Exercise, and Weight Loss Intervention Trial (DEW-IT). *Hypertension*. 2002 Nov;40(5):612-8.
- Mitrou PN, Kipnis V, Thiébaud AC, Reedy J, Subar AF, Wirfält E, Flood A, Mouw T, Hollenbeck AR, Leitzmann MF, Schatzkin A. Mediterranean dietary pattern and prediction of all-cause mortality in a US population: results from the NIH-AARP Diet and Health Study. *Arch Intern Med*. 2007 Dec 10;167(22):2461-8.

- Nowson CA, Wattanapenpaiboon N, Pachett A. Low-sodium Dietary Approaches to Stop Hypertension-type diet including lean red meat lowers blood pressure in postmenopausal women. *Nutr Res.* 2009 Jan;29(1):8-18.
- Nowson CA, Worsley A, Margerison C, Jorna MK, Frame AG, Torres SJ, Godfrey SJ. Blood pressure response to dietary modifications in free-living individuals. *J Nutr.* 2004 Sep;134(9):2322-9.
- Nowson CA, Worsley A, Margerison C, Jorna MK, Godfrey SJ, Booth A. Blood pressure change with weight loss is affected by diet type in men. *Am J Clin Nutr.* 2005 May;81(5):983-9.
- Núñez-Córdoba JM, Valencia-Serrano F, Toledo E, Alonso A, Martínez-González MA. The Mediterranean diet and incidence of hypertension: the Seguimiento Universidad de Navarra (SUN) Study. *Am J Epidemiol.* 2009 Feb 1;169(3):339-46.
- Osler M, Heitmann BL, Gerdes LU, Jørgensen LM, Schroll M. Dietary patterns and mortality in Danish men and women: a prospective observational study. *Br J Nutr.* 2001 Feb;85(2):219-25.
- Panagiotakos D, Pitsavos C, Chrysohoou C, Palliou K, Lentzas I, Skoumas I, Stefanadis C. Dietary patterns and 5-year incidence of cardiovascular disease: a multivariate analysis of the ATTICA study. *Nutr Metab Cardiovasc Dis.* 2009 May;19(4):253-63.
- Panagiotakos DB, Pitsavos C, Matalas AL, Chrysohoou C, Stefanadis C. Geographical influences on the association between adherence to the Mediterranean diet and the prevalence of acute coronary syndromes, in Greece: the CARDIO2000 study. *Int J Cardiol.* 2005 Apr 8;100(1):135-42.
- Parikh A, Lipsitz SR, Natarajan S. Association between a DASH-like diet and mortality in adults with hypertension: findings from a population-based follow-up study. *Am J Hypertens.* 2009 Apr;22(4):409-16.
- Rouse IL, Beilin LJ, Armstrong BK, Vandongen R. Blood-pressure-lowering effect of a vegetarian diet: controlled trial in normotensive subjects. *Lancet.* 1983 Jan 1;1(8314-5):5-10.
- Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, Obarzanek E, Conlin PR, Miller ER 3rd, Simons-Morton DG, Karanja N, Lin PH; DASH-Sodium Collaborative Research Group. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med.* 2001 Jan 4;344(1):3-10.
- Schulze MB, Hoffmann K, Kroke A, Boeing H. Risk of hypertension among women in the EPIC-Potsdam Study: comparison of relative risk estimates for exploratory and hypothesis-oriented dietary patterns. *Am J Epidemiol.* 2003 Aug 15;158(4):365-73.
- Sciarrone SE, Strahan MT, Beilin LJ, Burke V, Rogers P, Rouse IR. Ambulatory blood pressure and heart rate responses to vegetarian meals. *J Hypertens.* 1993 Mar;11(3):277-85.
- Shimazu T, Kuriyama S, Hozawa A, Ohmori K, Sato Y, Nakaya N, Nishino Y, Tsubono Y, Tsuji I. Dietary patterns and cardiovascular disease mortality in Japan: a prospective cohort study. *Int J Epidemiol.* 2007 Jun;36(3):600-9.

Singman HS, Berman SN, Cowell C, Maslansky E, Archer M. The Anti-Coronary Club: 1957 to 1972. *Am J Clin Nutr*. 1980 Jun;33(6):1183-91.

Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: meta-analysis. *BMJ*. 2008 Sep 11;337:a1344.

Swain JF, McCarron PB, Hamilton EF, Sacks FM, Appel LJ. Characteristics of the diet patterns tested in the optimal macronutrient intake trial to prevent heart disease (OmniHeart): options for a heart-healthy diet. *J Am Diet Assoc*. 2008 Feb;108(2):257-65.

Takahashi Y, Sasaki S, Okubo S, Hayashi M, Tsugane S. Blood pressure change in a free-living population-based dietary modification study in Japan. *J Hypertens*. 2006 Mar;24(3):451-8.

Trichopoulou A, Bamia C, Trichopoulos D. Anatomy of health effects of Mediterranean diet: Greek EPIC prospective cohort study. *BMJ*. 2009 Jun 23;338:b2337.

Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med*. 2003 Jun 26;348(26):2599-608.

Trichopoulou A, Kouris-Blazos A, Vassilakou T, Gnardellis C, Polychronopoulos E, Venizelos M, Ligiou P, Wahlqvist ML, Trichopoulos D. Diet and survival of elderly Greeks: a link to the past. *Am J Clin Nutr*. 1995 Jun;61(6 Suppl):1346S-1350S.

Waijers PM, Ocké MC, van Rossum CT, Peeters PH, Bamia C, Chloptsios Y, van der Schouw YT, Slimani N, Bueno-de-Mesquita HB. Dietary patterns and survival in older Dutch women. *Am J Clin Nutr*. 2006 May;83(5):1170-6.

Willcox BJ, Willcox DC, Todoriki H, Fujiyoshi A, Yano K, He Q, Curb JD, Suzuki M. Caloric restriction, the traditional Okinawan diet, and healthy aging: the diet of the world's longest-lived people and its potential impact on morbidity and life span. *Ann N Y Acad Sci*. 2007 Oct;1114:434-55.

World Cancer Research Fund and American Institute for Cancer Research Report (WCRF/AICR). *Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective*. Washington, DC: AICR, 2007.

Table B2.5. Selected dietary patterns with documented cardiovascular health benefits (adjusted to 2000 calories)

Dietary Pattern	DASH with Reduced Sodium	Mediterranean Diet (Greece)	Mediterranean Diet (Spain)	Mediterranean Diet (US)	Japanese	Okinawan
Citation	Karanja et al, 1999 and Lin et al, 2003	Trichopoulou et al, NEJM 2003	Nunez-Cordoba 2008 (SUN Study; MAI high score)	Fung et al, 2009	Wilcox et al, 2007 (Circa 1950)	Wilcox et al, 2007 (Circa 1949)
Qualitative Description						
Emphasizes	Potassium-rich vegetables, fruits, and low-fat dairy products	Plant- foods, vegetables, fruits, grains, beans, nuts and seeds, olive oil, and fish	Plant- foods, vegetables, fruits, breads, other cereals potatoes, beans, nuts and seeds, olive oil, and fish	Plant foods, vegetables, fruits, whole grains, legumes, fish	Rice, legumes, soy foods, vegetables, seaweed, and fish	Plant-foods, primarily Okinawan sweet potatoes, rice, legumes, soy foods, other vegetables, and nutrient rich foods of low energy density
Includes	Whole grains, poultry, fish, and nuts	Lean meat Red wine	Cheese, yogurt Red wine	Lean meat	Fruit Meat and eggs	
Limits (small amount)	Red meats, sweets, and sugar-containing beverages		Red meat Sweets	Potatoes	Milk products	Fruit Meat, eggs Milk products
Nutrients						
Calories (kcal)	2000	2000	2000	2000	2000	2000
Carbohydrates (% total kcal)	58%	nd	47%	39.1%	79%	85%
Protein (% total kcal)	18%	nd	18%	15.1%	13%	9%
Total Fat (% total kcal)	27%	~42.7 (summed)	33%	nd	8%	6%
Saturated Fat (% total kcal)	7%	13.1 %	10%	10% (Incl. trans)	2.0%	1.9%
Monounsaturated (% total kcal)	10%	22.7%	15 %	9.5%	2.3%	1.8%
Polyunsaturated (% total kcal)	8%	6.9%	5.1 %	nd	3.5%	2.4%
Cholesterol (mg)	143	nd	nd	nd	nd	nd
Fiber (g)	29	nd	29	20	22	26
Potassium (mg)	4371	nd	4589	nd	2623	5826
Sodium (mg)	1095	nd	2532	nd	2370	1269

Table B2.5 (continued). Selected dietary patterns with documented cardiovascular health benefits (adjusted to 2000 calories)

Dietary Pattern	DASH with Reduced Sodium	Mediterranean Diet (Greece)	Mediterranean Diet (Spain)	Mediterranean Diet (US)	Japanese	Okinawan
Food Groups						
Vegetables: total (c)	2.1	4.1	1.2	2.2	nd	nd
- Dark Green (c)	Nd	nd	nd	nd	<0.1 (seaweed)	<0.1 (sea weed)
- Legumes ² (c)	Nd	<0.1	0.4	0.3	0.3	0.5
- Red Orange (c)	Nd	nd	nd	nd	0.5 (Asian sweet potatoes)	6.6 (Asian sweet potatoes)
- Other Veg (c)	Nd	nd	nd	nd	1.3; + 0.3 (pickled veg)	0.9
- Starchy Veg (c)	Nd	0.6	nd	No potatoes	0.3 (other potatoes)	<0.1 (other potatoes)
Fruit & juices (c)	2.5	1.0 (fruit & nuts) 1.5 (juice & other bev)	1.3 (fruit & juice) 0.1 (dried fruit & nuts)	1.6	0.2 (papaya & tomato = veg)	<0.1 (papaya & tomato = veg)
Grains: total (oz)	7.3	5.4	2.0	nd	2.4; 1.7 (rice)	1.1; 0.9 (rice)
- Whole grains (oz)	3.9	nd	nd	1.6	nd	nd
Milk & milk products, Whole	0.7	1.0	0.8	nd	<0.1	<0.1
- Low-fat (c)	1.9	nd	1.3	nd	nd	nd
Animal Proteins:						
- Meat (oz)	1.4	3.5	3.6	2.4	0.4	0.1
- Poultry (oz)	1.7	nd	nd	nd	nd	nd
- Eggs (oz)	Nd	nd	1.9	nd	0.3	<0.1
- Fish (total) (oz)	1.4	0.8	2.4	1.5	2.1	0.6
-- Hi N3 (oz)	Nd	nd	nd	nd	nd	nd
-- Low N3 (oz)	Nd	nd	nd	nd	nd	nd
Plant Proteins:						
- Legumes (oz)	0.4	nd	0.4	nd	0.4 (Incl soy)	0.3 (Incl soy)
- Nuts & seeds (oz)	0.9	See fruit above.	See fruit above.	0.5	< 1 g	<0.1
- Soy products (oz)	Nd	nd		nd	See legumes.	See legumes.
Oils (g)	24.8	40.3 (olive oil)	19.0 (olive oil)	nd	nd	nd
Solid Fats (g)	Nd	nd	nd	nd	nd	nd
Added Sugar (g)	12	24.3	nd	nd	7.7	3.4
Alcohol (g)	Nd	7.9 ²	7.1 (red wine)	7.3	30.0 (flavors and alcohol)	7.8 (flavors and alcohol)

Table B2.6. Dietary patterns and blood pressure in adults

Author and Year	Study Type	Quality	Population/Location	Sig SBP Reduction	Sig DBP Reduction	Caveats
DASH	N = 12 (9 RCT, 3 prospective cohort)	12 Positive 2 Neutral		12 +	10 + 1 Ø 1 n/d	
Appel LJ et al., 2005	RCT (OmniHeart)	Positive	N = 164 adult with prehypertension or stage 1 hypertension US	+	+	Overall Between Diet Differences - SBP: Pro vs.Cho diet: P =0.002; Unsat Fat vs. Cho: P = 0.005 DBP: Pro vs.Cho diet: P <0.001; Unsat Fat vs. Cho: P = 0.02
Appel LJ et al., 1997	RCT	Positive	N = 459; 234 males; 225 females Normo and hypertensive subjects US	+	+	SBP: P < 0.001 DBP: Males P <0.001; Females P = 0.003
Appel LJ et al., 2003	RCT	Positive	N = 810 free living adults Normo and Hypertensive US	+	+	SBP and DBP: P <0.001
Azadbakht L et al., 2005	RCT	Neutral	N =116 subjects with metabolic syndrome BP ≥ 130/85 Iran	+	+	For both men and women P<0.001
Dauchet L et al., 2007	Longitudinal and cross-sectional analysis	Positive	N= 6,119 (2596 men, 3523 women); free living France	+	+	SBP: P <0.05 DBP: P < 0.01 Longitudinal results: DASH score: SBP: P<0.002; DBP: P<0.02
Forman JP et al., 2009	Prospective cohort study	Positive	N = 83,882 females; Nurse's Health Study II Normotensive US	+	+	Outcome in multivariate HR (95% CI) for incident HTN
Miller ER et al., 2002	RCT	Positive	N = 43 US	+	+	SBP, DBP: P <0.001
Nowson CA et al., 2009	RCT	Positive	N = 111 females (menopausal) Australia	+ + **	Ø +**	SBP: P = 0.38, 0.21** DBP: P = 0.61, 0.27** ** With HTN meds

Table B2.6 (continued). Dietary patterns and blood pressure in adults

Author and Year	Study Type	Quality	Population/Location	Sig SBP Reduction	Sig DBP Reduction	Caveats
DASH	N = 12 (9 RCT, 3 prospective cohort)	12 Positive 2 Neutral		12 +	10 + 1 ∅ 1 n/d	
Nowson CA et al., 2004	RCT	Positive	N = 94 males and females Australia	+	+	SBP: P = 0.001 DBP: P = 0.05
Sacks FM et al., 2001	RCT (cross-over)	Positive	N = 390 (males, females; black and white) US	+	n/d	SBP: P < 0.001
Schulze MB et al., 2003	Prospective cohort study	Positive	N = 8,552 females Normotensive Germany	+	+	HR (95% CI) for incident HTN
MEDITERRANEAN	N = 1 cohort	1 Positive		1+	1+	
Núñez-Córdoba JM et al., AJE 2009	Prospective cohort study (6 yr f/u)	Positive	N = 9,408 adults; 3,583 males, 5,825 females Spain	+	+	SBP: P = 0.01 DBP: P = 0.05
VEGETARIAN	N = 4 RCT	3 Positive 1 Neutral		4+	3 + 1 ∅	
Hakala P and Karvetti RL, 1989	RCT	Positive	N = 110 adults Finland	+	+	SBP: P = 0.05 DBP: P = 0.01
Margetts BM et al., 1986	RCT (cross-over)	Neutral	N = 58; 42 males, 16 females Untreated mild hypertensives Australia	+	∅	SBP: P , 0.05
Rouse IL et al., 1983	RCT (cross-over)	Positive	N = 59 males and females Australia	+	+	SBP, DBP: P < 0.01
Sciarrone SE et al., 1993	RCT	Positive	N = 21 males Australia	+	+	Ovo-lacto vegetarian
Japanese/Okinawan	N = 1 RCT	1 Positive		1+	1 ∅	
Takahashi Y 2006	RCT	Positive	N = 550 (202 males, 348 females) Japan	+	∅	SBP: P = 0.007 Japanese diet with ↑ Vitamin C, carotene, Fruits and vegetables ↓ Sodium intake

Table B2.7. Dietary patterns, cardiovascular disease (CVD), and mortality in adults

Author and Year/ Quality/ Study Type	Population/ Location	CVD	Mortality	Outcomes	Comments/Caveats
DASH and DASH Variations	N=10 1 RCT 9 Cohort				
Appel et al., 2005 Randomized, 3-period Crossover Trial Positive	N=164 (mean age = 53.6 yr; 45% women) Omni-Heart US	+	ND	Compared with baseline, all diets lowered estimated CHD risk. Compared with the high carbohydrate diet, estimated 10-yr CHD risk was lower and similar on the high protein and high unsaturated fat diets. Compared to high carbohydrate diet, high UFA diet decreased SBP; increased HDL-C; decreased TG, no change in LDL-C	Addresses total fat question : High UFA diet replaced 10% energy from CHO (total fat=37% E; 21% MUFA; 10% PUFA; 6% SFA). High UFA improved CHD risk, BP, and serum lipids, compared to high CHO (SFA constant).
Folsom et al., 2007 Prospective Cohort Study Neutral	N = 20,993, 55-69 yrs at baseline Iowa Women's Health Study Non-hypertensive	∅	∅	Incidence of hypertension inversely associated w/ degree of concordance with DASH diet (P for trend = 0.02), After adjustment for additional risk factors, little evidence that any endpoint assoc w/ DASH score	DASH diet concordance score calculated w/ baseline FFQ in 1986, subjects followed through 2002.
Fung et al., 2001 Prospective Cohort Study Positive	N = 69,017, 38 - 63 yrs at baseline Nurses' Health Study US	+	ND	Higher prudent-pattern score assoc w/ lower risk total CHD (RR Q5 vs Q1=0.61, 95%CI: 0.49-0.76, P for trend <0.001); after adjustment for BMI, smoking, caloric intake, supplemental use, hormone replacement therapy, and other coronary risk factors (RR=0.76, 95% CI: 0.60-0.98, P for trend = 0.03). Higher western-pattern score assoc w/ higher risk total MI after adjusting for age (RR Q5 versus Q1= 1.44, 95%CI: 1.16-1.78, P for trend <.001); remained sig. after multivariate adjustment (RR=1.46, 95%CI: 1.07-1.99).	12 y follow-up: 1984-1996 Baseline=1984 All FFQs using 1984 format (116 item)
Fung et al., 2008 Prospective Cohort Study Positive	N = 88,517, 34 - 59 yrs at baseline Nurses' Health Study US	+	+	RR of CHD across quintiles of DASH score = 1.0, 0.99, 0.86, 0.87 and 0.76 (95% CI: 0.67 - 0.85, P for trend <0.001) Magnitude of risk difference was similar for nonfatal MI and fatal CHD DASH score assoc w/ ↓ risk of stroke	24y follow-up: 1980-2004 Baseline=1980 Included data from older 1980 FFQ (61 item) and 1984 FFQ
Heidemann et al., 2008 Prospective Cohort Study Positive	N = 72,113 Nurses' Health Study US	+	+	Prudent pattern assoc w/ 28% lower risk of cardiovascular mortality and 17% lower risk of all-cause mortality, Western pattern assoc w/ 22% higher risk of cardiovascular mortality, 16% higher risk of cancer, and 21% higher risk of all-cause mortality.	18 y follow-up: 1984-2002 Baseline=1984 All FFQs using 1984 format (116 item)

Table B2.7 (continued). Dietary patterns, cardiovascular disease (CVD), and mortality in adults

Author and Year/ Quality/ Study Type	Population/ Location	CVD	Mortality	Outcomes	Comments/Caveats
DASH and DASH Variations	N=10 1 RCT 9 Cohort				
Hu et al., 2000 Prospective Cohort Study Positive	N=44,875 men, 40-75 y at baseline Health Professionals Follow-up Study	+	+	Two patterns explaining < 20% of the variance identified by factor analysis: Prudent and Western Higher Prudent score assoc w/ monotonic lower risk of CHD (RR across quintiles: 1.0, 0.84, 0.76, 0.71, 0.66 (95% CI: 0.54-0.80, P for trend < 0.0001 For fatal CHD after adjustment for age, smoking, BMI, and other CHD risk factors (RR across increasing quintiles: 1.0, 0.83, 0.78, 0.81, 0.70 (95%CI: 0.54, 0.91, P for trend=0.03 Higher Western score assoc w/ monotonic higher risk of CHD (RR across quintiles (P<0.0001) CHD RR (highest Prudent vs lowest Western) = 0.50 (95%CI: 0.34, 0.74).	8 y follow-up from 1986 Authors conclude dietary patterns derived from their FFQ predict CHD risk independent of other lifestyle factors.
Levitan et al., 2009 Prospective Cohort Study Neutral	36,019 women, 48-83 y at baseline Swedish Mammography Cohort	+	+	Top quartile of DASH score had 37% lower rate of heart failure (HF); rate ratios across quartiles = 1 (ref), 0.85 (95% CI: 0.66-1.11), 0.69 (95% CI: 0.54-0.88), and 0.63 (95% CI: 0.48-0.81), P for trend <0.001. Both HF-assoc hospitalization and death were determined	7 y follow-up; dietary intake only measured at baseline Hypertension was based on self-report.
Osler et al., 2001 Prospective Cohort Study Neutral	N= 5,872 (2,994 men, 2,878 women) Random equal-sized samples 30,40,50, 60-y at baseline Danish World Health Organization MONICA survey	+	+	Prudent pattern inversely assoc w/ all-cause (hazard ratios =0.63 in women =0.75 in men) and cardiovascular mortality Western pattern not associated w/ mortality	
Parikh et al., 2009 Prospective Cohort Study Neutral	N=5532 adults w/ hypertension NHANES III (1988-1994) US	+	+	DASH-like group had lower unadjusted mortality rates per 1,000 person-yrs for all-cause mortality (P=0.02), stroke mortality (P<0.001), and cancer mortality (P=0.05). DASH-like group, after adjusting for multiple confounders, assoc w/ lower mortality from all causes (HR=0.69, 95% CI 0.52-0.92, P=0.01) and stroke (HR=0.11, 95%CI 0.03-0.47, P=0.003). CVD mortality risk (HR=0.92, 95%CI 0.63-1.35, P=0.67), IHD (HR=0.77, 95%CI 0.47-1.14, P=0.28), and cancer (HR=0.51, 95%CI 0.23-1.10, P=0.09) not stat significant	8.2 person-years follow-up Secondary outcomes included specific causes of mortality CVD, ischemic heart disease, stroke, and cancer

Table B2.7 (continued). Dietary patterns, cardiovascular disease (CVD), and mortality in adults

Author and Year/ Quality/ Study Type	Population/ Location	CVD	Mortality	Outcomes	Comments/Caveats
DASH and DASH Variations	N=10 1 RCT 9 Cohort				
Singman et al., 1980 Prospective Cohort Study Neutral	N=1,113 men experimental and 467 men control US	+	ND	Prudent diet group in both age categories (40-49 y & 50-59 y) had lower CHD incidence rates	
MEDITERRANEAN	N=13 1 Index 1 Systematic Rev 1 Meta Analysis 9 Cohort 1 Case Control				
Alberti et al., 2008 Analysis of Mediterranean Adequacy Index (MAI) Neutral	5 data sets on 23 populations	ND	+	Inverse correlation between MAI and 25 y CHD death rate and total mortality	MAI: divide the sum of the percentages of dietary energy from food groups typical of a healthy reference Mediterranean diet, by the sum of the percentages of dietary energy of food groups that are not characteristic of a healthy reference Mediterranean diet
Fidanza et al., 2004 Prospective Cohort Study Neutral	N=12,763 men, 40-59 yrs at baseline US	+	+	The coefficient of linear correlation between the MAI and CHD death rates in the 16 cohorts was -0.72 (P=0.001)	MAI Index
Fung et al., 2009 Prospective Cohort Study Neutral	N = 76,522 , 38 - 63 yrs at baseline Nurses' Health Study US	+	+	Top aMed quintile ↓risk CHD and stroke: RR CHD = 0.71, 95% CI: 0.62-0.82, P for trend < 0.0001, RR stroke = 0.87, 95% CI: 0.73-1.02, P for trend = 0.03 CVD mortality ↓: top quintile RR=0.61, 95% CI:0.49-0.76, P for trend <0.0001	20 y follow-up: 1984-2004 Baseline=1984 All FQs using 1984 format
Harriss et al., 2007 Prospective Cohort Study Neutral	N= 40,653 (16,673 men, 23,908 women) Melbourne Collaborative Study	+	+	Mediterranean dietary factor inversely assoc w/ CVD and IHD mortality IHD, HR (highest compared w/ lowest quartile) = 0.59 (95% CI: 0.39-0.89, P for trend=0.03) Excluding subjects w/ prior CVD (HR=0.51, 95% CI: 0.30-0.88, P for trend = 0.03)	Mean follow-up = 10.4 y Involved migrants to Australia from Mediterranean countries (24% of subjects were Mediterranean born)

Table B2.7 (continued). Dietary patterns, cardiovascular disease (CVD), and mortality in adults

Author and Year/ Quality/ Study Type	Population/ Location	CVD	Mortality	Outcomes	Comments/Caveats
MEDITERRANEAN	N=13 1 Index 1 Systematic Rev 1 Meta Analysis 9 Cohort 1 Case Control				
Knoops et al., 2004 Prospective Cohort Study Neutral	N= 40,653 (1,507 men, 832 women) HALE cohort Netherlands	+	+	Mediterranean diet (HR = 0.77, 95% CI: 0.68 - 0.88) assoc w/ ↓ risk all-cause mortality Similar results were observed for mortality from coronary heart disease, cardiovascular diseases, and cancer	10 y mortality from all causes (CVD, CHD, and Cancer)
Mente et al., 2009 Systematic Review/ Meta-analysis Positive	146 prospective cohort studies + 43 RCTs (pub1950-2007) Europe, Asia, US	+	ND	Among the dietary exposures with strong evidence of causation from cohort studies, only the Mediterranean dietary pattern is related to CHD in RCTs	Used Bradford Hill guidelines to derive causation score based on 4 criteria (strength, consistency, temporality, and coherence) for each dietary exposure in cohort studies and examined for consistency with the findings of RCTs.
Mitrou et al., 2007 Prospective Cohort Study Positive	N= 352,497 (196,158 men, 156,339 women) median age = 62 NIH-AARP Diet and Health Study US	+	+	Men: multivariate HR all-cause mortality = 0.79 (95%CI: 0.76 - 0.83), CVD mortality = 0.78 (95% CI: 0.69 - 0.87), cancer mortality = 0.83 (95% CI: 0.76 - 0.91). Women: ↓ risks = 12% cancer mortality (P for trend = 0.04); = 20% all-cause mortality (P for trend < 0.001).	5 y follow-up Used 9-point score to assess conformity with Mediterranean dietary pattern (components included vegetables, legumes, fruits, nuts, whole grains, fish, monounsaturated fat-saturated fat ratio, alcohol, and meat)
Panagiotakos et al., 2005 Case-control Study Positive	N= 848 w/ 1 st CHD event and 1,078 age- and sex-matched controls (aged 49 - 75) CARDIO2000 Study Greece	+	ND	10-unit increase in Mediterranean diet score assoc w/ 27% (95% CI: 0.66 - 0.89) decrease odds of non-fatal acute coronary syndromes	Secondary prevention
Panagiotakos et al., 2009 Prospective Cohort Study Neutral	N = 2,101 ATTICA Study Greece	+	ND	Pattern characterized by cereals, small fish, and olive oil assoc w/ ↓ CVD risk (HR = 0.72, 95% CI: 0.52 - 1.00) Pattern characterized by fruit and vegetables using olive oil in cooking (HR = 0.80, 95% CI: 0.66 - 0.97) Patterns characterized by sweets, red meat, margarine, salty nuts, hard cheese and alcohol assoc w/ ↑ CVD risk	5 y follow-up Exclusion of CVD done by detailed clinical evaluation

Table B2.7 (continued). Dietary patterns, cardiovascular disease (CVD), and mortality in adults

Author and Year/ Quality/ Study Type	Population/ Location	CVD	Mortality	Outcomes	Comments/Caveats
MEDITERRANEAN					
N=13 1 Index 1 Systematic Rev 1 Meta Analysis 9 Cohort 1 Case Control					
Trichopoulos et al., 2003 Prospective Cohort Study Neutral	N = 22,043, 38-63 yrs at baseline EPIC Study Greece	+	+	Higher adherence to Med diet assoc w/ ↓ total mortality (adjusted HR = 0.75, 95% CI: 0.64 - 0.87); inverse assoc w/ CHD death (adjusted HR = 0.67, 95% CI: 0.47 - 0.94) and cancer death (adjusted HR = 0.76, 95% CI: 0.59 - 0.98).	44 month follow-up
Trichopoulos et al., 2009 Prospective Cohort Study Neutral	N = 23,349 EPIC Study Greece	ND	+	Higher adherence to a Med diet assoc w/ ↓ total mortality (adjusted mortality ratio = 0.864, 95% CI: 0.802 - 0.932).	8.5 y follow-up
Waijers et al., 2006 Prospective Cohort Study Neutral	N = 5,427 women (aged >60 years) EPIC Study Netherlands	ND	+	Principal component analysis identified 3 diet patterns: Mediterranean, Traditional Dutch, and Healthy Dutch Healthy trad Dutch pattern assoc w/ ↓ mortality rate; women in highest tertile 30% ↓ mortality risk	8.2 y follow-up
VEGETARIAN					
N=5 4 Cohort 1 Time series					
Chang-Claude et al., 2005 Prospective Cohort Study Neutral	N = 1,904; 858 males, 1,046 females 1,165 lacto-ovo, 679 non-veg, 60 vegans. Germany	+	∅	↓ risk ischemic heart disease (RR = 0.70, 95% CI: 0.41 - 1.18) No effect on mortality (RR = 1.10, 95% CI: 0.89 - 1.36)	A cohort study of vegetarians and health-conscious persons in Germany was followed-up prospectively for 21 years, including 1,225 vegetarians and 679 health-conscious nonvegetarians
Fraser et al., 2005 Time series Neutral	(N=30,292 males, N=50,562 females) California Seventh Day Adventists (N=297,126 male, 344,401 female) Stanford Five-City Project US	+	+	Rate ratio (RR) (Adventist/Stanford study) 1 st event fatal CHD = 0.59 (95% CI, 0.43-0.80) men and 0.49 (0.32-0.76) women. Vegetarian Adventists, RR = 0.45 (0.24-0.84) and 0.20 (0.06-0.63) men and women, respectively. 1st event MI RR = 0.60 (0.47-0.78) and 0.46 (0.33-0.65). Vegetarian Adventists RR = 0.37 (0.20-0.66) and 0.62 (0.35-1.09) men and women, respectively.	Two concurrent California observational studies, one with unusual dietary habits, are compared. Similar diagnostic criteria were used in both the Adventist Health Study and the Stanford Five-City Project.

Table B2.7 (continued). Dietary patterns, cardiovascular disease (CVD), and mortality in adults

Author and Year/ Quality/ Study Type	Population/ Location	CVD	Mortality	Outcomes	Comments/Caveats
VEGETARIAN	N=5 4 Cohort 1 Time series				
Key et al., 1996 Prospective Cohort Study Neutral	N = 10,771; 4,336 males, 6,435 females UK	+	+	Daily consumption of fresh fruit assoc w/ ↓ mortality ischemic heart disease (rate ratio = 0.76, 95% CI: 0.60 - 0.97), cerebrovascular disease (rate ratio = 0.68, 95% CI: 0.47 - 0.98), and all causes (rate ratio = 0.79, 95% CI: 0.70 - 0.90)	Mortality ratios measured for vegetarianism and for daily consumption of wholemeal bread, bran cereals, nuts or dried fruit, fresh fruit, and raw salad in relation to all cause mortality and mortality from ischaemic heart disease, cerebrovascular disease, all malignant neoplasms, lung cancer, colorectal cancer, and breast cancer.
Key et al., 1998 Meta-analysis: 5 Prospective Cohort Studies Neutral	N = 76,172 men and women US	+	+	Compared to non-vegetarians, vegetarians had 24% ↓ IHD mortality (rate ratio = 0.76, 95% CI:0.62-0.94) Reduction in mortality among vegetarians varied significantly with age at death. Regular meat consumers compared to semi-vegetarians (fish or meat <1X/wk), IHD rate ratios=0.78 (95% CI:0.68-0.89) in semi-vegetarians and 0.66 (95% CI:0.53-0.83) in vegetarians (P for trend <0.001).	Vegetarians were those who did not eat any meat or fish (n = 27,808). Non-vegetarians were from a similar background to the vegetarians within each study.
Mann et al., 1997 Prospective Cohort Study Neutral	N = 10,802; 4,102 males, 6,700 females Health conscious, mean age=33-34 United Kingdom	+	+	An increase in mortality for IHD was observed with increasing intakes of total and saturated animal fat and dietary cholesterol- death rate ratios in the third tertile compared with the first tertile: 329, 95% confidence interval (CI) 150 to 721; 277, 95% CI 125 to 613; 353, 95% CI 157 to 796, respectively. No protective effects for dietary fiber, fish or alcohol	13.3 y follow-up Prospective observation of vegetarians, semi-vegetarians, and meat eaters
Japanese/Okinawan	N=1 Cohort				
Shimazu et al., 2007 Prospective Cohort Study Neutral	N=40,547, 40-79 yrs at baseline Japan	+	+	3 patterns identified by principal components analysis: i) a Japanese dietary pattern highly correlated with soybean products, fish, seaweeds, vegetables, fruits and green tea, (ii) an 'animal food' dietary pattern and (iii) a high-dairy, high-fruit-and-vegetable, low-alcohol (DFA) dietary pattern. Japanese pattern assoc w/ ↓ risk CVD mortality (HR = 0.73, 95% CI 0.59-0.92, P for trend=0.003)	7 y follow-up

ND = Not determined.

Part B. Section 3: Translating and Integrating the Evidence: A Call to Action

The data clearly document that America is experiencing a public health crisis involving overweight and obesity. Particularly alarming is the further evidence that the obesity epidemic involves American children and youth, as nearly one in three are classified as overweight or obese. Childhood obesity and overweight is a serious health concern in the United States (US) because of immediate health consequences, as well as because it places a child at increased risk of obesity in adulthood, with all its attendant health problems such as cardiovascular diseases (CVD) and type 2 diabetes (T2D). All adults—parents, educators, caregivers, teachers, policy makers, health care providers, and all other adults who work with and care about children and families—serve as role models in some capacity and share responsibility for helping the next generation prevent obesity by promoting healthy lifestyles at all ages. Primary prevention of obesity, starting in pregnancy and early childhood, is the single best strategy for combating and reversing America’s obesity epidemic for current and future generations. While there is also an urgent need to improve the health and well-being of children and adults who are already overweight and obese, primary prevention offers the strongest universal benefits. Solving the obesity problem will take a coordinated system-wide, multi-sectoral approach that engages parents as well as those in education, government, healthcare, agriculture, business, advocacy and the community. This approach must promote primary prevention among those who are not yet overweight and address weight loss and fitness among those who are overweight.

Disparities in health among racial and ethnic minorities and among different socioeconomic groups have been recognized as a significant concern for decades. Several subgroups of the population (Native Americans, Blacks, Hispanics, and segments of the population with low income) have a strikingly high prevalence of overweight and obesity. Dietary patterns vary among different ethnic and socioeconomic groups. Individuals of lower education and/or income levels tend to eat fewer servings of vegetables and fruits than do those with more education and/or higher income. According to national surveys, Blacks tend to have the lowest intakes of vegetables and fruits among ethnic groups, but also have a higher prevalence of hypertension and related diseases, such as stroke. Although the reasons for these differences are complex and multifactorial, this report addresses research indicating that certain dietary changes can provide a means to reduce health disparities. If we are successful in changing dietary intake patterns of all Americans through a systematic approach, we will go along way in narrowing the gap in health disparities.

Although obesity is related to many chronic health conditions, it is not the only diet-related public health problem confronting the nation. Nutritionally suboptimal diets with or without obesity are etiologically related to many of the most common, costly, and yet preventable health problems in

the US, particularly CVD (atherosclerosis, stroke) and related risk factors (T2D, hypertension, and hyperlipidemia), some cancers, and osteoporosis. Improved nutrition and appropriate eating behaviors have tremendous potential to enhance public health, prevent or reduce morbidity and mortality, and decrease health care costs.

The science is not perfect; evidence is strong in some areas and limited or inconsistent in other areas. Nevertheless, this report is an urgent call to action to address a major public health crisis by focusing on helping all Americans achieve energy balance through adoption and adherence to current nutrition and physical activity guidelines.

After reviewing its entire report, the Dietary Guidelines Advisory Committee (DGAC) recognized a need to not only document the evidence, but translate and integrate major findings that have cross-cutting public health impact and provide guidance on how to implement the changes necessary to enhance the health and well being of the population. Below are the four major cross-cutting findings from the 2010 DGAC Report, followed by suggestions for implementation.

Four Main Integrated Findings to be Used in Developing the 2010 Dietary Guidelines for Americans

1. Reduce the incidence and prevalence of overweight and obesity of the US population by reducing overall calorie intake and increasing physical activity.

A focus on life-stage approaches (pregnant women, children, adolescents, adults, and older adults) is necessary nationwide to help Americans meet nutrient needs within appropriate calorie intake. To achieve this, Americans should:

- Know their calorie needs. In other words, individuals need to know how many calories they should consume each day based on their age, sex, and level of physical activity.
- Significantly lower excessive calorie intake from added sugars, solid fats, and some refined grain products.
- Increase their consumption of a variety of vegetables, fruits, and fiber-rich whole grains.
- Avoid sugar-sweetened beverages.
- Consume smaller portions, especially of high-calorie foods.
- Choose lower-calorie options, especially when eating foods away from home.
- Increase their overall physical activity.

- Have access to improved, easy-to-understand labels listing calorie content and portion size on packaged foods and for restaurant meals (especially quick service [i.e., fast food] restaurants, restaurant chains, and other places where standardized foods are served).

Collectively, these measures will help Americans manage their body weight and improve their overall health. In order to achieve this goal, the public and private sectors must be committed to assisting all Americans to know their calorie needs at each stage of life and help them recognize how to manage and/or lower their body weight. Simple but effective consumer-friendly tools for self-assessment of energy needs and self-monitoring of food and beverage intake are urgently needed and should be developed. These strategies will enable everyone to recognize and implement, both inside and outside the home, dietary recommendations that have been consistent for decades.

2. Shift food intake patterns to a more plant-based diet that emphasizes vegetables, cooked dry beans and peas, fruits, whole grains, nuts, and seeds. In addition, increase the intake of seafood and fat-free and low-fat milk and milk products, and consume only moderate amounts of lean meats, poultry, and eggs.

This approach will help Americans meet their nutrient needs while maintaining energy balance. Importantly, this will assist Americans to increase their intake of shortfall nutrients, such as potassium and fiber. These goals can be attained through a range of food patterns—from omnivore to vegan—that embrace cultural heritage, lifestyle, and food preferences. These flexible patterns of eating must encompass all foods and beverages that are consumed as meals and snacks throughout the day, regardless of whether they are eaten at home or away from home.

3. Significantly reduce intake of foods containing added sugars and solid fats because these dietary components contribute excess calories and few, if any, nutrients. In addition, reduce sodium intake and lower intake of refined grains, especially refined grains that are coupled with added sugar, solid fat, and sodium.

The components of the American diet that are consumed in excess are solid fats and added sugars (SoFAS), refined grains, and sodium. SoFAS alone contribute approximately 35 percent to total energy intake of Americans. Collectively, the consumption of foods containing SoFAS, refined grains, and sodium lead to excessive calorie intake, resulting in weight gain and health consequences such as hypertension, CVD, and T2D. Reducing the intake of these over-consumed components will require much more than individual behavior change. A comprehensive approach is needed. The food industry will need to act to help Americans achieve these goals. Every aspect of the industry, from research and development to production and retail, needs to contribute healthful food

solutions to reduce the intake of SoFAS, certain refined grain products, and sodium. Sound health and wellness policies at the local, state, and national level also can help facilitate these changes.

4. Meet the 2008 Physical Activity Guidelines for Americans.

A comprehensive set of physical activity recommendations for people of all ages and physical conditions was released by the US Department of Health and Human Services in 2008 (HHS, 2008). The 2008 Physical Activity Guidelines for Americans were developed to help Americans to become more physically active. By objective measures, large portions, indeed the majority, of the US population are sedentary (Metzger, 2008). In fact, Americans spend most of their waking hours engaged in behaviors that expend very little energy (Matthews, 2008). To increase the public's participation in physical activity, compelling multi-sector approaches are needed to improve home, school, work, and community environments to promote physical activity. These changes need to surpass planned exercise and foster greater energy expenditure throughout the day. Improved exposure to recreational spaces, increased use of active transportation, and encouraging development of school and worksite policies that program physical activity throughout the day can help enable Americans to develop and maintain healthier lifestyle behaviors. Special attention and creative approaches also are needed to help Americans reduce sedentary behaviors, especially television viewing and video game use, among children and adolescents.

A Call to Action

Dietary Guidelines for Americans have been published since 1980. During this time obesity rates have escalated and dietary intake patterns have strayed from the ideal. The 2010 DGAC recognizes that several of its recommendations have been made repeatedly in prior reports with little or no demonstrable impact. For example, recommended intakes of vegetables and fruit remain woefully unchanged, despite continuing advice to markedly increase intake of these foods. Substantial, high-level barriers appear to impede achievement of these goals, including certain government regulations and policies. Chief among these are land use policy and economic incentives for food manufacturers. The food supply and access to it has changed dramatically over the past 40 years, contributing to an overall increased calorie intake by many individuals. Since the 1970s, the number of fast food restaurants has increased 147 percent. The portions that are served in restaurants and the serving sizes of foods sold in packages at stores have increased as well. Moreover, the number of food items at the supermarket has increased from 10,425 in 1978 to 46,852 in 2008, and most of these contribute SoFAS, refined grains, and sodium to the American diet (see *Part D. Section 1. Energy Balance and Weight Management* for a discussion of recent changes in the food environment). This has far-reaching effects such that the average child now

consumes 365 calories per day of added sugars and 433 calories per day of solid fat for a combined total of 798 calories, or more than one-third of total calorie intake (HHS, 2010; see *Part D. Section 2. Nutrient Adequacy*). Conversely, Americans spend 45 percent less time preparing food at home (see *Part D. Section 1. Energy Balance and Weight Management*) or eating food at the family table than previously, and this behavioral trend is associated with increased risk of weight gain, overweight and obesity. **In this context, the DGAC concluded that mere repetition of advice will not effectively help Americans achieve these evidence-based and often-repeated goals for a healthy diet.**

Ensuring that all Americans consume a health-promoting dietary pattern and achieve and maintain energy balance requires far more than individual behavior change. A multi-sectoral strategy is imperative. For this reason, the 2010 DGAC strongly recommends that HHS and USDA convene appropriate committees, potentially through the Institute of Medicine (IOM), to develop a strategic plan focusing on the behaviors and actions needed to successfully implement the four key 2010 DGAC recommendations highlighted above.

A coordinated strategic plan that includes all sectors of society, including individuals, families, educators, communities, allied health professionals, public health advocates, policy makers, scientists, and small and large businesses (e.g., farmers, agricultural producers, food scientists, food manufacturers, and food retailers of all kinds), should be engaged in developing and implementing the plan to help all Americans eat well, be physically active, and maintain good health. It is important that any strategic plan be evidence-informed, action-oriented, and focused on changes in systems (IOM, 2010a). This systems approach is already underway in countries such as the United Kingdom for obesity prevention (Butland, 2007) with promising results. Recent examples of this approach in the US include an IOM committee convened by HHS and USDA and charged with developing strategies for gradually but dramatically reducing sodium intake, which remains persistently high even after more than 40 years of advice. This IOM committee recently issued its report (IOM, 2010b), providing a comprehensive strategy to reduce dietary sodium intake in the general population by focusing on the food supply and targeting industry to partner in systematic reductions in sodium content of foods. Already there is encouraging evidence that food manufacturers are responding positively and are committed to reducing the sodium content in their food products. Similarly, the US National Physical Activity Plan, released in May, 2010, was developed by multiple stakeholders and provides a comprehensive, realistic implementation framework intended to promote physical activity in the American population. Most recently, the May, 2010, White House Task Force on Childhood Obesity Report, *Solving the Problem of Childhood Obesity Within a Generation*, also calls for a multi-sector, systems approach to solving this important public health issue.

An Urgent Need to Focus on Children

Any and all systems-based strategies must include a focus on children. Primary prevention of obesity must begin in childhood. This is the single most powerful public health approach to combating and reversing America's obesity epidemic over the long term. Trends for childhood overweight and obesity are alarming, with obesity prevalence rates tripling between 1980 and 2004. Although rates for children appear to be leveling off, they remain high, with one-third currently overweight or obese, defined as at or above the 85th percentile on body mass index (BMI)-for-age growth charts (Ogden, 2010). These numbers represent more than 25 million children in the US. In order to reverse this trend, we will need to work together as a Nation to improve the food environment to which children are exposed at home, school, and the community. Efforts to prevent childhood obesity need to start very early, even in utero. Increasing evidence indicates that maternal obesity before conception and excessive gestational weight gain represent a substantial risk of childhood obesity in the offspring (see *Part D. Section 2. Energy Balance and Weight Management* for a detailed discussion of this issue). Thus, addressing maternal nutrition, physical activity, and body weight before conception and during pregnancy as well as emphasizing early childhood nutrition is paramount for preventing the onset of childhood obesity. Areas targeting childhood obesity prevention that should be addressed include, but are not limited to:

- Improve foods sold and served in schools, including school breakfast, lunch, and after-school meals and competitive foods so that they meet the recommendations of the IOM report on school meals (IOM, 2009) and the key findings of the 2010 DGAC. This includes all age groups of children, from preschool through high school.
- Increase comprehensive health, nutrition, and physical education programs and curricula in US schools and preschools, including food preparation, food safety, cooking, and physical education classes and improved quality of recess.
- Develop nationally standardized approaches for health care providers to track BMI-for-age and provide guidance to children and their families to effectively prevent, monitor, and/or treat childhood obesity.
- Develop nationally standardized approaches for health care providers to improve nutrition, physical activity participation, healthy weight gain during pregnancy and the attainment of a healthy weight postpartum.
- Increase safe routes to schools and community recreational areas to encourage active transportation and physical activity.
- Remove sugar-sweetened beverages and high-calorie snacks from schools, recreation facilities, and other places where children gather.

- Develop and enforce responsible zoning policies for the location of fast food restaurants near schools and places where children play.
- Increase awareness and promote action around reducing screen time (television and computer or game modules) and removing televisions from children's bedrooms.
- Develop and enforce effective policies regarding marketing of food and beverage products to children. Efforts in this area are underway through a government interagency committee comprised of the Federal Trade Commission, Centers for Disease Control and Prevention, USDA, and Food and Drug Administration, as well as some self-regulation from industry (Omnibus Appropriations Act, 2009).
- Develop affordable summer programs that support children's health, as children gain the most weight during the out-of-school summer months (von Hippel, 2007).

Challenges and Opportunities for Change

Change is needed in the overall food environment to support the efforts of all Americans to meet the key recommendations of the 2010 DGAC (Story, 2009). The 2010 DGAC recognizes that the current food environment does not adequately facilitate the ability of Americans to follow the evidence-based recommendations outlined in the 2010 DGAC Report. Population growth, availability of fresh water, arable land constraints, climate change, current policies, and business practices are among some of the major challenges that need to be addressed in order to ensure that these recommendations can be implemented nationally. For example, if every American were to meet the vegetable, fruit, and whole-grain recommendations, domestic crop acreage would need to increase by an estimated 7.4 million harvested acres (Buzby, 2006). Furthermore, the environment does not facilitate the ability of individuals to follow the 2008 Physical Activity Guidelines for Americans. Most home, school, work, and community environments do not promote engagement in a physically active lifestyle. To meet these challenges, the following sustainable changes must occur:

- Improve nutrition literacy and cooking skills, and empower and motivate the population to prepare and consume healthy foods at home, especially among families with children.
- For all Americans, especially those with low-income, create greater financial incentives to purchase, prepare, and consume vegetables and fruit, whole grains, seafood, fat-free and low-fat milk and milk products, lean meats, and other healthy foods. Currently, individuals have an economic disincentive to purchase healthy foods.
- Improve the availability of affordable fresh produce through greater access to grocery stores, produce trucks, and farmers' markets.

- Increase environmentally sustainable production of vegetables, fruits, and fiber-rich whole grains.
- Ensure household food security through measures that provide access to adequate amounts of foods that are nutritious and safe to eat.
- Develop safe, effective, and sustainable practices to expand aquaculture and increase the availability of seafood to all segments of the population. Ensure that consumers have access to user-friendly benefit/risk information to make informed seafood choices.
- Encourage restaurants and the food industry to offer health-promoting foods that are low in sodium; limited in SoFAS and refined grains; and served in smaller portions.
- Implement the US National Physical Activity Plan, a private-public sector collaborative promoting local, state, and national programs and policies to increase physical activity and reduce sedentary activity (National Physical Activity Plan, 2010). Through the Plan and other initiatives, develop efforts across all sectors of society, including health care and public health; education; business and industry; mass media; parks, recreation, fitness, and sports; transportation, land use and community design; and volunteer and non-profit. Reducing screen time, especially television, for all Americans also will be important.

The 2010 DGAC recognizes the significant challenges involved in implementing the goals outlined here. These challenges go beyond cost, economic interests, technological and societal changes, and agricultural limitations. Over the past several decades, the value of preparing and enjoying healthy food has eroded, leaving instead the practices of eating processed foods containing excessive sodium, solid fats, refined grains, and added sugars. As a Nation, we all need to value and adopt the practices of good nutrition, physical activity, and a healthy lifestyle. The DGAC encourages all stakeholders to take actions to make every choice available to Americans a healthy choice. To move toward this vision, all segments of society—from parents to policy makers and everyone else in between—must now take responsibility and play a leadership role in creating gradual and steady change to help current and future generations live healthy and productive lives. A measure of success will be evidence that meaningful change has occurred when the 2015 DGAC convenes.

References

- Butland B, Jebb S, Kopelman P, McPherson K, Thomas S, Mardell J, Parry V. *Foresight. Tackling Obesity: Future Choice—Project Report*. London (UK): Government Office for Science; 2007. <http://www.foresight.gov.uk/Obesity/17.pdf>. Accessed May 5, 2010.
- Buzby JC, Wells HF, Vocke G. *Possible Implications for U.S. Agriculture from Adoption of Select Dietary Guidelines*. Washington, DC: US Department of Agriculture, Economic Research Service Report No. (ERR-31); 2006.
- Institute of Medicine (IOM). *School Meals: Building Blocks for Healthy Children*. Washington, DC: The National Academies Press; 2009.
- Institute of Medicine (IOM). *Bridging the Evidence Gap in Obesity Prevention: A Framework to Inform Decision Making*. Washington, DC: The National Academies Press; 2010a.
- Institute of Medicine (IOM). *Strategies to Reduce Sodium Intake in the United States*. Washington, DC: The National Academies Press; 2010b.
- Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, Troiano RP. Amount of Time Spent in Sedentary Behaviors in the United States, 2003–2004. *Am J Epidemiol* 2008;167(7):875-81.
- Metzger JS, Catellier DJ, Evenson KR, Treuth MS, Rosamond WD, Siega-Riz AM. Patterns of objectively measured physical activity in the United States. *Med Sci Sports Exerc* 2008;40(4):630-38.
- Ogden CL, Carroll MD, Curtin LR, Lamb MM, Flegal KM. Prevalence of high body mass index in US children and adolescents, 2007-2008. *JAMA* 2010;303(3):242-9.
- Omnibus Appropriations Act, Pub. L. No. 111-8. 11 March 2009. Print.
- Story M, Hamm MW, Wallinga D. Food systems and public health: linkages to achieve healthier diets and healthier communities. *J Hunger and Environ Nut* 2009;4(3):219-24.
- National Physical Activity Plan. The US National Physical Activity Plan. <http://www.physicalactivityplan.org/theplan.htm>. Updated May 5, 2010. Accessed May 5, 2010.
- US Department of Health and Human Services (HHS). *2008 Physical Activity Guidelines for Americans*. Washington, DC: US Department of Health and Human Services. ODPHP Publication No. U0036; 2008. <http://www.health.gov/paguidelines>.
- US Department of Health and Human Services (HHS). Sources of selected nutrients among the US population, 2005-06. Risk Factor Monitoring and Methods Branch Website. Applied Research Program. National Cancer Institute. <http://riskfactor.cancer.gov/diet/foodsources/>. Accessed May 5, 2010.

von Hippel PT, Powell B, Downey DB, Rowland NJ. The effect of school on overweight in childhood: gain in body mass index during the school year and during summer vacation. *Am J Public Health*. 2007;97(4):696–702.